



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/437, 31/4184, 31/4439, C07D 471/04, 401/12, 235/28	A1	(11) International Publication Number: WO 00/50037 (43) International Publication Date: 31 August 2000 (31.08.00)
(21) International Application Number: PCT/US00/02524 (22) International Filing Date: 25 February 2000 (25.02.00) (30) Priority Data: 60/122,111 26 February 1999 (26.02.99) US (71) Applicant (for all designated States except US): NITROMED, INC. [US/US]; 12 Oak Park Drive, Bedford, MA 01730 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): GARVEY, David, S. [US/US]; 10 Grand Hill Drive, Dover, MA 02030 (US). LETTS, L., Gordon [AU/US]; 12 Abbott Road, Dover, MA 02030 (US). TAM, Sang, William [US/US]; 148 Pine Street, Dover, MA 02030 (US). WANG, Tiansheng [CN/US]; 2 Dumbur Way, Concord, MA 01742 (US). RICHARDSON, Stewart, K. [GB/US]; 55 Autumn Drive, Tolland, CT 06084 (US). (74) Agent: GRIEFF, Edward, D.; Hale and Dorr LLP, 1455 Pennsylvania Avenue, N.W., Washington, D.C. 20004 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NITROSATED AND NITROSYLATED PROTON PUMP INHIBITORS, COMPOSITIONS AND METHODS OF USE (57) Abstract <p>The present invention describes novel nitrosated and/or nitrosylated proton pump inhibitor compounds, and novel compositions comprising at least one proton pump inhibitor compound that is optionally substituted with at least one NO and/or NO₂ group, and, optionally, at least one compound that donates, transfers or releases nitric oxide, stimulates endogenous synthesis of nitric oxide, elevates endogenous levels of endothelium-derived relaxing factor or is a substrate for nitric oxide synthase, and/or at least one nonsteroidal antiinflammatory drug, selective COX-2 inhibitor antacid, bismuth-containing reagent, acid-degradable antibacterial compound, and mixtures thereof. The present invention also provides methods for treating and/or preventing gastrointestinal disorders; facilitating ulcer healing; decreasing the recurrence of ulcers; improving gastroprotective properties, anti-<i>Helicobacter pylori</i> properties or antacid properties of proton pump inhibitors; decreasing or reducing the gastrointestinal toxicity associated with the use of nonsteroidal antiinflammatory compounds; treating <i>Helicobacter pylori</i> and viral infections. The compounds and/or compositions of the present invention can also be provided in the form of a pharmaceutical kit.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**NITROSATED AND NITROSYLATED PROTON PUMP INHIBITORS,
COMPOSITIONS AND METHODS OF USE
RELATED APPLICATIONS**

This application claims priority to U. S. Provisional Application No.
5 60/122,111 filed February 26, 1999.

FIELD OF THE INVENTION

The present invention describes novel nitrosated and/or nitrosylated
proton pump inhibitor compounds, and novel compositions comprising at least
one proton pump inhibitor compound that is optionally substituted with at least
10 one NO and/or NO₂ group, and, optionally, at least one compound that donates,
transfers or releases nitric oxide, stimulates endogenous synthesis of nitric oxide,
elevates endogenous levels of endothelium-derived relaxing factor or is a
substrate for nitric oxide synthase, and/or at least one nonsteroidal
antiinflammatory drug, selective COX-2 inhibitor, antacid, bismuth-containing
15 reagent, acid-degradable antibacterial compound, and mixtures thereof. The
present invention also provides methods for treating and/or preventing
gastrointestinal disorders; facilitating ulcer healing; decreasing the recurrence of
ulcers; improving gastroprotective properties, anti-*Helicobacter pylori* properties
or antacid properties of proton pump inhibitors; decreasing or reducing the
20 gastrointestinal toxicity associated with the use of nonsteroidal antiinflammatory
compounds; treating *Helicobacter pylori* and viral infections. The compounds
and/or compositions of the present invention can also be provided in the form
of a pharmaceutical kit.

BACKGROUND OF THE INVENTION

25 The proton pump, located in the apical membrane of the parietal cell, is
responsible for the secretion of acid in the stomach when it is stimulated by the
enzyme adenosine triphosphate (H⁺, K⁺)-ATPase. Proton pump inhibitors are a
class of anti-secretory compounds used in the management of gastrointestinal
disorders. They suppress gastric acid secretion by the specific inhibition of the
30 (H⁺, K⁺)-ATPase enzyme system at the secretory surface of the gastric parietal cell.

A family of substituted benzimidazoles have been developed as specific
proton pump inhibitors. Two of these compounds, omeprazole and
lansoprazole, are used clinically in the United States. Structurally they contain a

sulfinyl group bridging between substituted benzimidazole and pyridine rings. At a neutral pH, omeprazole and lansoprazole are chemically stable, are weak bases, are lipid-soluble, and do not show any inhibitory activity. Once these compounds reach the parietal cells and diffuse into the secretory canaliculi, they become protonated. The protonated compounds rearrange to form sulfenic acid and then a sulfenamide. The latter interacts covalently with sulfhydryl groups at critical sites in the extracellular (luminal) domain of the membrane spanning (H⁺, K⁺)-ATPase. Inhibition occurs when two molecules of the inhibitor are bound per molecule of the enzyme. The specificity of these proton pump inhibitors arises from the selective distribution of the (H⁺, K⁺)-ATPase, the acid-catalyzed rearrangement of the compounds to generate the active inhibitor, and the trapping of the protonated compound and the cationic sulfenamide within the acidic canaliculi and adjacent to the target enzyme.

Omeprazole and lansoprazole are typically administered orally as delay-release capsules. The compounds are stable at a neutral pH, but are destroyed by gastric acid. Therefore, if the integrity of the gelatin-coated capsule is destroyed in any way and the patient swallows the enteric-coated grains, the neutral pH in the mouth and the esophagus will break down the microencapsulation, and the compounds will be degraded by the gastric acid in the stomach. The delay release capsules, when appropriately taken, release the omeprazole and lansoprazole after the granules leave the stomach.

Despite their good anti-secretory properties, proton pump inhibitors are not unanimously recognized as gastroprotective agents. In addition, there is a high relapse rate associated with treating gastrointestinal disorders with proton pump inhibitors as they do not eliminate *Helicobacter pylori* (*Campylobacter pylori*), the bacteria responsible for peptic ulcer disease, gastric lymphoma and adenocarcinoma. U.S. Patent Nos. 5,599,794 and 5,629,305 describe the administration of proton pump inhibitors in combination with acid-degradable antibacterial compounds for the treatment of infections caused by *Helicobacter pylori*.

A variety of adverse reactions have been ascribed to proton pump inhibitors, such as omeprazole and lansoprazole, reflecting, in part, the very large number of patients who have been treated with these drugs. The incidence of

adverse reactions is low, and the adverse reactions are generally minor. Due to the profound reduction in gastric acidity, there tends to be an increased secretion of gastrin. Hence, patients who take therapeutic doses of omeprazole and lansoprazole have modest hypergastrinemia. Prolonged administration of high
5 doses of the drugs can cause hyperplasia of oxyntic mucosal cells.

The most common side effects of proton pump inhibitors, such as omeprazole and lansoprazole, are nausea, diarrhea, abdominal colic, and central nervous system effects such as headaches and dizziness. Occasionally skin rashes and transient elevations of plasma activities of hepatic aminotransferase have
10 been reported. The drugs can also result in bacterial overgrowth in the gastrointestinal tract and the development of nosocomial pneumonia.

There is a need in the art for proton pump inhibitors that have improved gastroprotective properties, decrease the recurrence of ulcers, facilitate ulcer healing and that can be used at low dosages. The present invention is directed to
15 these, as well as other, important ends.

SUMMARY OF THE INVENTION

The present invention provides proton pump inhibitors to which is linked at least one NO and/or NO₂ group (i.e., nitrosylated and/or nitrosated). The proton pump inhibitors can be, for example, substituted benzimidazoles and
20 substituted azabenzimidazoles, including, for example, omeprazole, pantoprazole, rabeprazole, leminoprazole, lansoprazole, timoprazole, tenatoprazole, disulprazole, esomeprazole, RO 18-5362 and IY 81149. The present invention also provides compositions comprising such compounds in a pharmaceutically acceptable carrier.

Another aspect of the invention provides compositions comprising at least one proton pump inhibitor, that is optionally substituted with at least one NO and/or NO₂ group (i.e., nitrosylated and/or nitrosated), and at least one compound that donates, transfers or releases nitric oxide and/or stimulates
25 endogenous production of nitric oxide (NO) or endothelium-derived relaxing factor (EDRF) *in vivo* and/or is a substrate for nitric oxide synthase. The present
30 invention also provides compositions comprising such compounds in a pharmaceutically acceptable carrier.

Yet another aspect of the present invention provides methods for treating

gastrointestinal disorders, methods for improving the gastroprotective properties, anti-*Helicobacter* properties and antacid properties of proton pump inhibitors, methods for facilitating ulcer healing and methods for decreasing the rate of recurrence of ulcers in a patient in need thereof comprising administering
5 to the patient at least one proton pump inhibitor that is substituted with at least one NO and/or NO₂ group, or by administering to the patient at least one proton pump inhibitor, that is optionally substituted with at least one NO and/or NO₂ group, and at least one compound that donates, transfers or releases nitric oxide and/or stimulates endogenous production of nitric oxide or EDRF *in vivo*. The
10 proton pump inhibitor, that is optionally substituted with at least one NO and/or NO₂ group, and nitric oxide donor can be administered separately or as components of the same composition in one or more pharmaceutically acceptable carriers.

The present invention also provides methods to decrease or reverse
15 gastrointestinal toxicity resulting from the administration of nonsteroidal antiinflammatory drugs (NSAIDs) and/or selective COX-2 inhibitors, and methods to facilitate ulcer healing resulting from the administration of NSAIDs and/or selective COX-2 inhibitors, in a patient in need thereof by administering the compounds and/or compositions of the present invention. The present
20 invention also provides methods to treat infections caused by *Helicobacter pylori* and/or viruses in patients in need thereof by administering the compounds and/or compositions of the present invention.

These and other aspects of the present invention are described in detail herein.

25 BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows the gastric lesion scores of (a) vehicle alone (open bar, n=10); (b) lansoprazole in vehicle (stripped bar, n=10); and (c) example 2 (nitrosylated lansoprazole) in vehicle (checked bar, n=9). Lansoprazole at 200 µmol/kg did not significantly inhibit the formation of gastric lesions relative to vehicle alone.
30 Example 2 (nitrosylated lansoprazole) at 200 µmol/kg inhibited the formation of gastric lesions relative to vehicle alone (p<0.05) and lansoprazole (p<0.05).

DETAILED DESCRIPTION OF THE INVENTION

As used throughout the disclosure, the following terms, unless otherwise

indicated, shall be understood to have the following meanings.

"Gastrointestinal disorder" refers to any disease or disorder of the upper and lower gastrointestinal tract of a patient including, for example, inflammatory bowel disease, Crohn's disease, irritable bowel syndrome, ulcerative colitis, peptic
5 ulcers, stress ulcers, bleeding peptic ulcers, duodenal ulcers, infectious enteritis, colitis, diverticulitis, gastric hyperacidity, dyspepsia, gastroparesis, Zollinger-Ellison syndrome, gastroesophageal reflux disease, *Helicobacter Pylori* associated disease, short-bowel (anastomosis) syndrome, hypersecretory states associated with systemic mastocytosis or basophilic leukemia and hyperhistaminemia that
10 result, for example, from neurosurgery, head injury, severe body trauma or burns.

"Upper gastrointestinal tract" refers to the esophagus, the stomach, the duodenum and the jejunum.

"Lower gastrointestinal tract" refers to the ileum, the colon, the cecum and
15 the rectum.

"Ulcers" refers to lesions of the upper gastrointestinal tract lining that are characterized by loss of tissue. Such ulcers include gastric ulcers, duodenal ulcers and gastritis.

"Viral infection" refers to both RNA and DNA viral infections. The RNA
20 viral infections include, but are not limited to, orthomyxoviridae, paramyxoviridae, picornaviridae, rhabdoviridae, coronavaridae, togaviridae, bunyaviridae, arenaviridae and reteroviridae. The DNA viral infections include, but are not limited to, adenoviridae, proxviridae, papovaviridae, herpesviridae and herpesviridae. The most preferable viral infections are those of the
25 herpesviridae family, such as, for example, herpes simplex viruses HSV-1 and HSV-2, cytomegalovirus (CMV), herpes varicella-zoster (VZV), Epstein-Barr (EBV), HHV6, HHV7, pseudorabies and rhinotracheitis, and the like.

"Proton pump inhibitor" refers to any compound that reversibly or irreversibly blocks gastric acid secretion by inhibiting the H^+/K^+ -ATP ase enzyme
30 system at the secretory surface of the gastric parietal cell.

"NSAID" refers to a nonsteroidal anti-inflammatory compound or a nonsteroidal anti-inflammatory drug. NSAIDs inhibit cyclooxygenase, the enzyme responsible for the biosyntheses of the prostaglandins and certain

autocoid inhibitors, including inhibitors of the various isozymes of cyclooxygenase (including but not limited to cyclooxygenase-1 and -2), and as inhibitors of both cyclooxygenase and lipoxygenase.

5 "Cyclooxygenase-2 (COX-2) inhibitor" refers to a compound that selectively inhibits the cyclooxygenase-2 enzyme over the cyclooxygenase-1 enzyme. Preferably, the compound has a cyclooxygenase-2 IC_{50} of less than about 0.5 μM , and also has a selectivity ratio of cyclooxygenase-2 inhibition over cyclooxygenase-1 inhibition of at least 50, and more preferably of at least 100. Even more preferably, the compound has a cyclooxygenase-1 IC_{50} of greater than
10 about 1 μM , and more preferably of greater than 20 μM . The compound can also inhibit the enzyme, lipoxygenase.

"Patient" refers to animals, preferably mammals, more preferably humans.

"Transdermal" refers to the delivery of a compound by passage through the skin and into the blood stream.

15 "Transmucosal" refers to delivery of a compound by passage of the compound through the mucosal tissue and into the blood stream.

"Penetration enhancement" or "permeation enhancement" refers to an increase in the permeability of the skin or mucosal tissue to a selected pharmacologically active compound such that the rate at which the compound
20 permeates through the skin or mucosal tissue is increased.

"Carriers" or "vehicles" refers to carrier materials suitable for compound administration and include any such material known in the art such as, for example, any liquid, gel, solvent, liquid diluent, solubilizer, or the like, which is non-toxic and which does not interact with any components of the composition
25 in a deleterious manner.

"Nitric oxide adduct" or "NO adduct" refers to compounds and functional groups which, under physiological conditions, can donate, release and/or directly or indirectly transfer any of the three redox forms of nitrogen monoxide (NO^+ , NO^- , NO^\bullet), such that the biological activity of the nitrogen monoxide species is
30 expressed at the intended site of action.

"Nitric oxide releasing" or "nitric oxide donating" refers to methods of donating, releasing and/or directly or indirectly transferring any of the three redox forms of nitrogen monoxide (NO^+ , NO^- , NO^\bullet), such that the biological

activity of the nitrogen monoxide species is expressed at the intended site of action.

"Nitric oxide donor" or "NO donor" refers to compounds that donate, release and/or directly or indirectly transfer a nitrogen monoxide species, and/or stimulate the endogenous production of nitric oxide or endothelium-derived relaxing factor (EDRF) *in vivo* and/or elevate endogenous levels of nitric oxide or EDRF *in vivo*. "NO donor" also includes compounds that are substrates for nitric oxide synthase.

"Alkyl" refers to a lower alkyl group, a haloalkyl group, an alkenyl group, an alkynyl group, a bridged cycloalkyl group, a cycloalkyl group or a heterocyclic ring, as defined herein.

"Lower alkyl" refers to branched or straight chain acyclic alkyl group comprising one to about ten carbon atoms (preferably one to about eight carbon atoms, more preferably one to about six carbon atoms). Exemplary lower alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, pentyl, neopentyl, iso-amyl, hexyl, octyl, and the like.

"Haloalkyl" refers to a lower alkyl group, an alkenyl group, an alkynyl group, a bridged cycloalkyl group, a cycloalkyl group or a heterocyclic ring, as defined herein, to which is appended one or more halogens, as defined herein. Exemplary haloalkyl groups include trifluoromethyl, chloromethyl, 2-bromobutyl, 1-bromo-2-chloro-pentyl, and the like.

"Alkenyl" refers to a branched or straight chain C_2 - C_{10} hydrocarbon (preferably a C_2 - C_8 hydrocarbon, more preferably a C_2 - C_6 hydrocarbon) which can comprise one or more carbon-carbon double bonds. Exemplary alkenyl groups include propylenyl, buten-1-yl, isobutenyl, penten-1-yl, 2,2-methylbuten-1-yl, 3-methylbuten-1-yl, hexan-1-yl, hepten-1-yl, octen-1-yl, and the like.

"Alkynyl" refers to an unsaturated acyclic C_2 - C_{10} hydrocarbon (preferably a C_2 - C_8 hydrocarbon, more preferably a C_2 - C_6 hydrocarbon) which can comprise one or more carbon-carbon triple bonds. Exemplary alkynyl groups include ethynyl, propynyl, butyn-1-yl, butyn-2-yl, pentyl-1-yl, pentyl-2-yl, 3-methylbutyn-1-yl, hexyl-1-yl, hexyl-2-yl, hexyl-3-yl, 3,3-dimethyl-butyn-1-yl, and the like.

"Bridged cycloalkyl" refers to two or more cycloalkyl groups, heterocyclic groups, or a combination thereof fused via adjacent or non-adjacent atoms.

Bridged cycloalkyl groups can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, hydroxy, halo, carboxyl, alkylcarboxylic acid, aryl, amidyl, ester, alkylcarboxylic ester, carboxamido, alkylcarboxamido, oxo and nitro. Exemplary
5 bridged cycloalkyl groups include adamantyl, decahydronaphthyl, quinuclidyl, 2,6-dioxabicyclo[3.3.0]octane, 7-oxabicyclo[2.2.1]heptyl, 8-azabicyclo[3,2,1]oct-2-enyl and the like.

"Cycloalkyl" refers to a saturated or unsaturated cyclic hydrocarbon comprising from about 3 to about 8 carbon atoms. Cycloalkyl groups can be
10 unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, arylamino, diarylamino, alkylaryl amino, aryl, amidyl, ester, hydroxy, halo, carboxyl, alkylcarboxylic acid, alkylcarboxylic ester, carboxamido, alkylcarboxamido, oxo and nitro. Exemplary cycloalkyl groups include cyclopropyl, cyclobutyl,
15 cyclopentyl, cyclohexyl, cyclohexenyl, cyclohepta,1,3-dienyl, and the like.

"Heterocyclic ring or group" refers to a saturated, unsaturated cyclic or polycyclic hydrocarbon group having about 3 to about 12 carbon atoms (preferably about 4 to about 6 carbon atoms) where 1 to about 4 carbon atoms are replaced by one or more nitrogen, oxygen and/or sulfur atoms. Sulfur maybe in the thio,
20 sulfinyl or sulfonyl oxidation state. The heterocyclic ring or group can be fused to an aromatic hydrocarbon group. Heterocyclic groups can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, arylamino, diarylamino, alkylaryl amino, hydroxy, oxo, thial, halo, carboxyl, carboxylic ester,
25 alkylcarboxylic acid, alkylcarboxylic ester, aryl, arylcarboxylic acid, arylcarboxylic ester, amidyl, ester, carboxamido, alkylcarboxamido, arylcarboxamido, sulfonic acid, sulfonic ester, sulfonamido and nitro. Exemplary heterocyclic groups include pyrrolyl, 3-pyrrolyl, 4,5,6-trihydro-2H-pyran, pyridinyl, 1,4-dihydropyridinyl, pyrazolyl, triazolyl, pyrimidinyl, pyridazinyl, oxazolyl,
30 thiazolyl, imidazolyl, indolyl, thiophenyl, furanyl, tetrahydrofuran, tetrazolyl, 2-pyrrolyl, 3-pyrrolyl, pyrrolidinyl, oxazolindinyl, 1,3-dioxolanyl, 2,6-dioxabicyclo[3,3,0]octanyl, 2-imidazolinyl, imidazolindinyl, 2-pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,3-triazolyl, 1,3,4-

thiadiazolyl, 2H-pyranyl, 4H-pyranyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl, 1,3,5-trithianyl, benzo(b)thiophenyl, benzimidazolyl, quinolinyl, and the like.

"Heterocyclic compounds" refer to mono- and polycyclic compounds comprising at least one aryl or heterocyclic ring.

"Aryl" refers to a monocyclic, bicyclic, carbocyclic or heterocyclic ring system comprising one or two aromatic rings. Exemplary aryl groups include phenyl, pyridyl, naphthyl, quinoyl, tetrahydronaphthyl, furanyl, indanyl, indenyl, indoyl, and the like. Aryl groups (including bicyclic aryl groups) can be unsubstituted or substituted with one, two or three substituents independently selected from alkyl, alkoxy, amino, alkylamino, dialkylamino, arylamino, diarylamino, alkylarylamino, hydroxy, carboxyl, carboxylic ester, alkylcarboxylic acid, alkylcarboxylic ester, aryl, arylcarboxylic acid, arylcarboxylic ester, alkylcarbonyl, arylcarbonyl, amidyl, ester, carboxamido, alkylcarboxamido, carbomyl, sulfonic acid, sulfonic ester, sulfonamido and nitro. Exemplary substituted aryl groups include tetrafluorophenyl, pentafluorophenyl, sulfonamide, alkylsulfonyl, arylsulfonyl, and the like.

"Alkylaryl" refers to an alkyl group, as defined herein, to which is appended an aryl group, as defined herein. Exemplary alkylaryl groups include benzyl, phenylethyl, hydroxybenzyl, fluorobenzyl, fluorophenylethyl, and the like.

"Arylalkyl" refers to an aryl radical, as defined herein, attached to an alkyl radical, as defined herein.

"Cycloalkylalkyl" refers to a cycloalkyl radical, as defined herein, attached to an alkyl radical, as defined herein.

"Heterocyclicalkyl" refers to a heterocyclic ring radical, as defined herein, attached to an alkyl radical, as defined herein.

"Cycloalkenyl" refers to an unsaturated cyclic hydrocarbon having about 3 to about 10 carbon atoms (preferably about 3 to about 8 carbon atoms, more preferably about 3 to about 6 carbon atoms) comprising one or more carbon-carbon double bonds.

"Arylheterocyclic ring" refers to a bi- or tricyclic ring comprised of an aryl ring, as defined herein, appended via two adjacent carbon atoms of the aryl ring

to a heterocyclic ring, as defined herein. Exemplary arylheterocyclic rings include dihydroindole, 1,2,3,4-tetra-hydroquinoline, and the like.

"Alkoxy" refers to $R_{50}O-$, wherein R_{50} is an alkyl group, as defined herein. Exemplary alkoxy groups include methoxy, ethoxy, t-butoxy, cyclopentyloxy, and the like.

"Arylalkoxy or alkoxyaryl" refers to an alkoxy group, as defined herein, to which is appended an aryl group, as defined herein. Exemplary arylalkoxy groups include benzyloxy, phenylethoxy, chlorophenylethoxy, and the like.

"Alkoxyalkyl" refers to an alkoxy group, as defined herein, appended to an alkyl group, as defined herein. Exemplary alkoxyalkyl groups include methoxymethyl, methoxyethyl, isopropoxymethyl, and the like.

"Alkoxyhaloalkyl" refers to an alkoxy group, as defined herein, appended to a haloalkyl group, as defined herein. Exemplary alkoxyhaloalkyl groups include 4- methoxy-2-chlorobutyl and the like.

"Cycloalkoxy" refers to $R_{54}O-$, wherein R_{54} is a cycloalkyl group or a bridged cycloalkyl group, as defined herein. Exemplary cycloalkoxy groups include cyclopropyloxy, cyclopentyloxy, cyclohexyloxy, and the like.

"Haloalkoxy" refers to a haloalkyl group, as defined herein, to which is appended an alkoxy group, as defined herein. Exemplary haloalkyl groups include 1,1,1-trichloroethoxy, 2-bromobutoxy, and the like.

"Hydroxy" refers to $-OH$.

"Oxo " refers to $=O$.

"Oxy " refers to $-O^+R_{77}$, wherein R_{77} is an organic or inorganic cation.

"Organic cation" refers to a positively charged organic ion. Exemplary organic cations include alkyl substituted ammonium cations, and the like.

"Inorganic cation" refers to a positively charged metal ion. Exemplary inorganic cations include Group I metal cations such as for example, sodium, potassium, and the like.

"Hydroxyalkyl" refers to a hydroxy group, as defined herein, appended to an alkyl group, as defined herein.

"Amino" refers to $-NH_2$.

"Nitrate" refers to $-O-NO_2$.

"Nitrite" refers to $-O-NO$.

"Thionitrate" refers to $-S-NO_2$.

"Thionitrite" and "nitrosothiol" refer to $-S-NO$.

"Nitro" refers to the group $-NO_2$ and "nitrosated" refers to compounds that have been substituted therewith.

5 "Nitroso" refers to the group $-NO$ and "nitrosylated" refers to compounds that have been substituted therewith.

"Nitrile" and "cyano" refer to $-CN$.

"Halogen" or "halo" refers to iodine (I), bromine (Br), chlorine (Cl), and/or fluorine (F).

10 "Alkylamino" refers to $R_{50}NH-$, wherein R_{50} is an alkyl group, as defined herein. Exemplary alkylamino groups include methylamino, ethylamino, butylamino, cyclohexylamino, and the like.

"Arylamino" refers to $R_{55}NH-$, wherein R_{55} is an aryl group, as defined herein.

15 "Dialkylamino" refers to $R_{50}R_{52}N-$, wherein R_{50} and R_{52} are each independently an alkyl group, as defined herein. Exemplary dialkylamino groups include dimethylamino, diethylamino, methyl propargylamino, and the like.

"Diarylamino" refers to $R_{55}R_{60}N-$, wherein R_{55} and R_{60} are each independently an aryl group, as defined herein.

20 "Alkylarylamino" refers to $R_{50}R_{55}N-$, wherein R_{50} is an alkyl group, as defined herein, and R_{55} is an aryl group, as defined herein.

"Aminoalkyl" refers to an amino group, an alkylamino group, a dialkylamino group, an arylamino group, a diarylamino group, an alkylarylamino group or a heterocyclic ring, as defined herein, to which is
25 appended an alkyl group, as defined herein.

"Aminoaryl" refers to an amino group, an alkylamino group, a dialkylamino group, an arylamino group, a diarylamino group, an alkylarylamino group or a heterocyclic ring, as defined herein, to which is
30 appended an aryl group, as defined herein.

"Thio" refers to $-S-$.

"Sulfinyl" refers to $-S(O)-$.

"Methanthial" refers to $-C(S)-$.

"Thial" refers to =S.

"Sulfonyl" refers to $-S(O)_2$.

"Sulfonic acid" refers to $-S(O)_2OR_{76}$, wherein R_{76} is a hydrogen, an organic cation or an inorganic cation.

5 "Alkylsulfonic acid" refers to a sulfonic acid group, as defined herein, appended to an alkyl group, as defined herein.

"Arylsulfonic acid" refers to a sulfonic acid group, as defined herein, appended to an aryl group, as defined herein

10 "Sulfonic ester" refers to $-S(O)_2OR_{58}$, wherein R_{58} is an alkyl group, an aryl group, an alkylaryl group or an aryl heterocyclic ring, as defined herein.

"Sulfonamido" refers to $-S(O)_2-N(R_{51})(R_{57})$, wherein R_{51} and R_{57} are each independently a hydrogen atom, an alkyl group, an aryl group, an alkylaryl group, or an arylheterocyclic ring, as defined herein, and R_{51} and R_{57} when taken together are a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group,
15 as defined herein.

"Alkylsulfonamido" refers to a sulfonamido group, as defined herein, appended to an alkyl group, as defined herein.

"Arylsulfonamido" refers to a sulfonamido group, as defined herein, appended to an aryl group, as defined herein.

20 "Sulfamoyl" refers to $R_{51}S(O)_2-N(R_{57})-$, wherein R_{51} and R_{57} are each independently a hydrogen atom, an alkyl group, an aryl group, an alkylaryl group, or an arylheterocyclic ring, as defined herein, and R_{51} and R_{57} when taken together are a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group, as defined herein.

25 "Alkylthio" refers to $R_{50}S-$, wherein R_{50} is an alkyl group, as defined herein.

"Arylthio" refers to $R_{55}S-$, wherein R_{55} is an aryl group, as defined herein.

30 "Cycloalkylthio" refers to $R_{54}S-$, wherein R_{54} is a cycloalkyl group or a bridged cycloalkyl group, as defined herein. Exemplary cycloalkylthio groups include cyclopropylthio, cyclopentylthio, cyclohexylthio, and the like.

"Alkylsulfinyl" refers to $R_{50}-S(O)-$, wherein R_{50} is an alkyl group, as defined herein.

"Alkylsulfonyl" refers to $R_{50}-S(O)_2-$, wherein R_{50} is an alkyl group, as

defined herein.

"Arylsulfinyl" refers to $R_{55}-S(O)-$, wherein R_{55} is an aryl group, as defined herein.

"Arylsulfonyl" refers to $R_{55}-S(O)_2-$, wherein R_{55} is an aryl group, as defined
5 herein.

"Amidyl" refers to $R_{51}C(O)N(R_{57})-$ wherein R_{51} and R_{57} are each independently a hydrogen atom, an alkyl group, an aryl group, an alkylaryl group, or an arylheterocyclic ring, as defined herein.

"Ester" refers to $R_{51}C(O)O-$ wherein R_{51} is a hydrogen atom, an alkyl group, an aryl group, an alkylaryl group, or an arylheterocyclic ring, as defined herein.
10

"Carbamoyl" refers to $-O-C(O)N(R_{51})(R_{57})$, wherein R_{51} and R_{57} are each independently a hydrogen atom, an alkyl group, an aryl group, an alkylaryl group or an arylheterocyclic ring, as defined herein, or R_{51} and R_{57} taken together are a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group, as defined
15 herein.

"Carboxyl" refers to $-C(O)OR_{76}$, wherein R_{76} is a hydrogen, an organic cation or an inorganic cation, as defined herein.

"Carbonyl" refers to $-C(O)-$.

"Alkylcarbonyl" or "alkanoyl" refers to $R_{50}-C(O)-$, wherein R_{50} is an alkyl group, as defined herein.
20

"Arylcarbonyl" or "aroyl" refers to $R_{55}-C(O)-$, wherein R_{55} is an aryl group, as defined herein.

"Carboxylic ester" refers to $-C(O)OR_{58}$, wherein R_{58} is an alkyl group, an aryl group, an alkylaryl group or an aryl heterocyclic ring, as defined herein.

"Alkylcarboxylic acid" and "alkylcarboxyl" refer to an alkyl group, as
25 defined herein, appended to a carboxyl group, as defined herein.

"Alkylcarboxylic ester" refers to an alkyl group, as defined herein, appended to a carboxylic ester group, as defined herein.

"Arylcarboxylic acid" refers to an aryl group, as defined herein, appended
30 to a carboxyl group, as defined herein.

"Arylcarboxylic ester" and "arylcarboxyl" refer to an aryl group, as defined herein, appended to a carboxylic ester group, as defined herein.

"Carboxamido" refers to $-C(O)N(R_{51})(R_{57})$, wherein R_{51} and R_{57} are each

independently a hydrogen atom, an alkyl group, an aryl group, an alkylaryl group or an arylheterocyclic ring, as defined herein, and R_{51} and R_{57} when taken together are a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group, as defined herein.

5 "Alkylcarboxamido" refers to an alkyl group, as defined herein, appended to a carboxamido group, as defined herein.

"Arylcarboxamido" refers to an aryl group, as defined herein, appended to a carboxamido group, as defined herein.

10 "Urea" refers to $-N(R_{59})-C(O)N(R_{51})(R_{57})$ wherein R_{51} , R_{57} , and R_{59} are each independently a hydrogen atom, an alkyl group, an aryl group, an alkylaryl group, or an arylheterocyclic ring, as defined herein, or R_{51} and R_{57} taken together are a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group, as defined herein.

15 "Phosphoryl" refers to $-P(R_{70})(R_{71})(R_{72})$, wherein R_{70} is a lone pair of electrons, sulfur or oxygen, and R_{71} and R_{72} are each independently a covalent bond, a hydrogen, a lower alkyl, an alkoxy, an alkylamino, a hydroxy or an aryl, as defined herein.

20 Compounds that donate, transfer or release nitric oxide species *in vivo* have a wide spectrum of advantages and applications. The present invention is based on the discovery of the effects of such compounds together with one or more proton pump inhibitors and/or one or more proton pump inhibitors directly or indirectly linked with one or more nitric oxide moieties. Treatment or prevention of gastrointestinal disorders, improved gastroprotective properties, decreased rate of recurrence of peptic ulcers and faster ulcer healing can be
25 obtained by the use of the nitrosated and/or nitrosylated proton pump inhibitors of the present invention. Treatment or prevention of gastrointestinal disorders, improved gastroprotective properties, decreased rate of recurrence of peptic ulcers and faster ulcer healing can also be obtained by the use of one or more proton pump inhibitors, that are optionally directly or indirectly linked with one
30 or more nitric oxide moieties, in conjunction with one or more compounds that donate, release or transfer nitric oxide or stimulate endogenous production of nitric oxide or EDRF *in vivo* or are substrates for nitric oxide synthase.

Proton pump inhibitors are compounds that selectively inhibit gastric acid

secretion by the specific inhibition of the (H⁺, K⁺)-ATPase enzyme system at the secretory surface of the gastric parietal cell. A nitric oxide donor is a compound that contains at least one nitric oxide adduct and releases or chemically transfers a biologically active nitrogen monoxide species.

5 The compounds and compositions of the present invention are novel and can be used to treat numerous gastrointestinal diseases and disorders. Such gastrointestinal disorders include, for example, inflammatory bowel disease, Crohn's disease, irritable bowel syndrome, ulcerative colitis, peptic ulcers, stress ulcers, bleeding peptic ulcers, duodenal ulcers, infectious enteritis, colitis, 10 diverticulitis, gastric hyperacidity, gastric hyperacidity, dyspepsia, gastroparesis, Zollinger-Ellison syndrome, gastroesophageal reflux disease, *Helicobacter Pylori* associated disease, short-bowel (anastomosis) syndrome, hypersecretory states associated with systemic mastocytosis or basophilic leukemia and hyperhistaminemia that result, for example, from neurosurgery, head injury, 15 severe body trauma or burns. The compounds and compositions of the present invention can also be used as a pre-anesthetic medication in emergency operations to reduce the danger of aspiration of acidic gastric contents

 The proton pump inhibitors used in the compounds and compositions of the present invention can be any of those known in the art, such as those 20 exemplified herein.

 Omeprazole, i.e., 5-methoxy-2-((4-methoxy-3,5-dimethyl-2-pyridinyl)methyl)sulfinyl)-1H-benzimidazole, (marketed under the trade name PRILOSEC® by Astra Merck, Wayne, PA) and lansoprazole, i.e., 2-(((3-methyl-4-(2,2,2-trifluoro-ethoxy)-2-pyridinyl)methyl)sulfinyl)-1H-benzimidazole (marketed 25 under the trade name PREVACID® by TAP Pharmaceutical Inc., Deerfield, IL) are two of the most widely used compounds that inhibit gastric acid secretion. Other useful compounds include rabeprazole, i.e., 2-(((4-(3-methoxypropoxy)-3-methyl-2-pyridinyl)methyl)sulfinyl)-1H-benzimidazole (marketed under the trade name ACIPHEX® by Eisai, Inc.), pantoprazole, i.e., 5-(difluoromethoxy)-2-(((3,4- 30 dimethoxy-2-pyridinyl)methyl)sulfinyl)-1H-benzimidazole, leminoprazole, timoprazole, tenatoprazole, disulprazole, esomeprazole, RO 18-5362, IY 81149. These compound do not exhibit anticholinergic or histamine H₂-receptor antagonist properties, but suppress gastric acid secretion by the specific inhibition

of (H⁺, K⁺)-ATPase enzyme system at the secretory surface of the gastric parietal cell. As this enzyme system is regarded as the acid (proton) pump within the parietal cell, these substituted benzimidazoles have been characterized as gastric-acid-pump inhibitors as they block the final step of acid production. Although
5 the proton pump inhibitor anti-secretory agents are effective in treating gastrointestinal disorders, they do not have any gastroprotective properties and, in addition, there is a high recurrence of ulcers associated with their use.

Another group of proton pump inhibitors are substituted quinolines, which include, for example, 3-butyl-4-(2-methylphenylamino)-8-(2-
10 hydroxyethoxy)-quinoline.


Other proton pump inhibitors are disclosed in, for example, U. S. Patent Nos. 4,045,564, 4,255,431, 4,634,710, 4,758,579, 4,806,549, 4,806,550, 4,818,760, 4,839,365, 4,845,118, 4,871,734, 4,873,337, 4,956,366, 4,981,861, 5,114,955, 5,149,702, 5,439,917, 5,554,631, 5,665,730, 5,677,302, 5,686,458, 5,703,097, 5,750,531, 5,990,311,
15 5,952,504 and 5,945,425 and in EP 0 033 094 B1, EP 0 045 200 A1, EP 0 221 041 A2, EP 0 234 485 A1, EP 0 246 774 A1, EP 0 254 588 A1, EP 0 259 174 A1 and in WO 89/08104, WO 92/12969, WO 95/27714, WO 97/32854, WO 98/18784, WO 98/43968, WO 98/54172; the disclosures of each of which are incorporated by reference herein in their entirety.

Several of the above contemplated proton pump inhibitors are described more fully in the literature, such as in Goodman and Gilman, The Pharmacological Basis of Therapeutics (9th Edition), McGraw-Hill, pages 901-915 (1996); Merck Index on CD-ROM, Twelfth Edition, Version 12:1, (1996); STN Express, file phar and file registry, the disclosures of which are incorporated by
25 reference herein in their entirety.

The proton pump inhibitors of the present invention can be nitrosated and/or nitrosylated through one or more sites such as oxygen (hydroxyl condensation), sulfur (sulfhydryl condensation), carbon and nitrogen. The proton pump inhibitor compounds that are nitrosated and/or nitrosylated in
30 accordance with the invention and/or are included in the compositions of the invention can be any of those known in the art, including those exemplified below.

In one embodiment, the present invention describes nitrosated and/or

nitrosylated proton pump inhibitors of Formula (I) or a pharmaceutically acceptable salt thereof:



(I)

wherein

A is S, S(O), or S(O)₂;

B is -CNR₇R₇' or nitrogen;


J is CH or nitrogen;

R₁ is a hydrogen, an alkoxy group, a lower alkyl group, or an alkylthio group;

R₂ is a hydrogen, an alkoxy group, a lower alkyl group, an alkylthio group, a haloalkoxy group, an alkoxyalkyl group, -NR₇R₇', -OD₁, or -SD₁; or R₂ and R₁ taken together with the carbon chain to which they are attached form a cycloalkyl ring or a heterocyclic ring; or R₂ and R₃ taken together with the carbon chain to which they are attached form a cycloalkyl ring or a heterocyclic ring;

R₃ and R₁₁ are each independently a hydrogen, an alkoxy group, a lower alkyl group, or an alkylthio group; or R₃ and R₁₁ taken together with the carbon chain to which they are attached form a cycloalkyl ring or a heterocyclic ring;

R₄ and R₅ are each independently a hydrogen, an alkyl group, a halo group, an alkoxy group, a haloalkyl group, a haloalkoxy group, a cyano group, an aryl group, a heterocyclic ring, -NR₇R₇', -OD₁, or -CO₂R₁₂; or R₄ and R₅ taken together are:



wherein

R_6 is oxygen or $N=O-R_7$;

5 R_7 and R_7' are each independently hydrogen, a lower alkyl group or D; or R_7 and R_7' taken together with the nitrogen to which they are attached form a heterocyclic ring;

R_{10} is a hydrogen; or R_{10} and R_1 taken together with the carbon chain to which they are attached form a cycloalkyl ring;

10 R_{12} is a lower alkyl group or D;

D_1 is a hydrogen or D;

D is Q or K;

Q is $-NO$ or $-NO_2$;

K is $-W_a-E_b-(C(R_e)(R_f))_p-E_c-(C(R_e)(R_f))_x-W_d-(C(R_e)(R_f))_y-W_i-E_j-W_g-(C(R_e)(R_f))_z-$

15 T-Q;

a, b, c, d, g, i and j are each independently an integer from 0 to 3;

p, x, y and z are each independently an integer from 0 to 10;

W at each occurrence is independently $-C(O)-$, $-C(S)-$, $-T-$, $-(C(R_e)(R_f))_h-$, an alkyl group, an aryl group, a heterocyclic ring, an arylheterocyclic ring, or

20 $-(CH_2CH_2O)_q-$;

E at each occurrence is independently $-T-$, an alkyl group, an aryl group, $-(C(R_e)(R_f))_h-$, a heterocyclic ring, an arylheterocyclic ring, or $-(CH_2CH_2O)_q-$;

h is an integer from 1 to 10;

q is an integer from 1 to 5;

25 R_e and R_f are each independently a hydrogen, an alkyl, a cycloalkoxy, a halogen, a hydroxy, an hydroxyalkyl, an alkoxyalkyl, an arylheterocyclic ring, an alkylaryl, a cycloalkylalkyl, a heterocycloalkyl, an alkoxy, a haloalkoxy, an amino, an alkylamino, a dialkylamino, an arylamino, a diarylamino, an alkylaryl amino, an alkoxyhaloalkyl, a haloalkoxy, a sulfonic acid, a sulfonic ester, an alkylsulfonic

acid, an arylsulfonic acid, an arylalkoxy, an alkylthio, an arylthio, a cycloalkylthio, a cycloalkenyl, a cyano, an aminoalkyl, an aminoaryl, an aryl, an arylalkyl, an alkylaryl, a carboxamido, a alkylcarboxamido, an arylcarboxamido, an amidyl, a carboxyl, a carbamoyl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarbonyl, an arylcarbonyl, an ester, a carboxylic ester, an alkylcarboxylic ester, an arylcarboxylic ester, a haloalkoxy, a sulfonamido, an alkylsulfonamido, an arylsulfonamido, a sulfonic ester, a urea, a phosphoryl, a nitro, -T-Q, or $(C(R_e)(R_f))_k$ -T-Q, or R_e and R_f taken together with the carbons to which they are attached form a carbonyl, a methanthial, a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group;

k is an integer from 1 to 3;

T at each occurrence is independently a covalent bond, a carbonyl, an oxygen, $-S(O)_o-$ or $-N(R_a)R_i-$;

o is an integer from 0 to 2;

R_a is a lone pair of electrons, a hydrogen or an alkyl group;

R_i is a hydrogen, an alkyl, an aryl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarboxylic ester, an arylcarboxylic ester, an alkylcarboxamido, an arylcarboxamido, an alkylaryl, an alkylsulfinyl, an alkylsulfonyl, an arylsulfinyl, an arylsulfonyl, a sulfonamido, a carboxamido, a carboxylic ester, an amino alkyl, an amino aryl, $-CH_2-C(T-Q)(R_e)(R_f)$, or $-(N_2O_2)^- \cdot M^+$, wherein M^+ is an organic or inorganic cation; with the proviso that when R_i is $-CH_2-C(T-Q)(R_e)(R_f)$ or $-(N_2O_2)^- \cdot M^+$, or R_e or R_f are T-Q or $(C(R_e)(R_f))_k$ -T-Q, then the "-T-Q" subgroup can be a hydrogen, an alkyl, an alkoxy, an alkoxyalkyl, an aminoalkyl, a hydroxy, a heterocyclic ring or an aryl group; and


with the proviso that the compounds of Formula (I) must contain at least one nitrite, nitrate, thionitrite or thionitrate group.

In cases where R_e and R_f are a heterocyclic ring or taken together R_e and R_f are a heterocyclic ring, then R_i can be a substituent on any disubstituted nitrogen contained within the radical where R_i is as defined herein.

In cases where multiple designations of variables which reside in sequence are chosen as a "covalent bond" or the integer chosen is 0, the intent is to denote a single covalent bond connecting one radical to another. For example, E_0 would

denote a covalent bond, while E_2 denotes (E-E) and $(C(R_e)(R_f))_2$ denotes $-C(R_e)(R_f)-C(R_e)(R_f)-$.

Another embodiment of the present invention describes compounds of Formula (II) or pharmaceutically acceptable salts thereof:



II

wherein

R_8 is a lower alkyl group, an alkoxyalkyl group, an alkylaryl group, a cycloalkyl group, a cycloalkylalkyl group, an aryl group, an alkylaryl group, or K;


R_9 at each occurrence is independently a hydrogen, a lower alkyl group, an alkylthio group, a halogen, a cyano group, an alkanoyl group, a haloalkyl group, a carbamoyl group, $-NR_7D_1$, $-OD_1$, or $-CO_2R_{12}$;

R_{71} is a hydrogen, a lower alkyl group, an alkoxy group, or $-OD_1$;

J, K, D_1 , R_7 , R_{12} , q and o are as defined herein; and

with the proviso that the compounds of Formula (II) must contain at least one nitrite, nitrate, thionitrite or thionitrate group.

Another embodiment of the present invention describes compounds of Formula (III) or pharmaceutically acceptable salts thereof:



III


wherein

R_{13} and R_{14} are each independently a hydrogen, a lower alkyl group, an alkoxyalkyl, or a lower alkyl-OD₁; or R_{13} and R_{14} taken together along with the carbons to which they are attached form a cycloalkyl group or an aryl group;


R_{17} is a hydrogen or a lower alkyl group;

Y_3 is:

(a)



(b)



or

wherein

R_{15} is a hydrogen or a lower alkyl group;

R_{16} is a hydrogen, a halogen, or a lower alkyl group;

R_{63} is a lower alkyl group or a phenyl group;

A_1 , A_2 and A_3 comprise the other subunits of a 5- or 6-membered monocyclic aromatic ring and A_1 , A_2 and A_3 are each independently:


(i) CR_o , wherein R_o at each occurrence is hydrogen or -OD₁;

- (ii) $N-R_p$, wherein R_p at each occurrence is independently a covalent bond to an adjacent ring atom in order to render the ring aromatic, a hydrogen, or K;
- (iii) a sulfur atom;
- (iv) an oxygen atom; or
- (v) $B_a = B_b$, wherein B_a and B_b are each independently a nitrogen atom or CR_o ; wherein R_o at each occurrence is hydrogen or $-OD_1$;

D_1 and K are as defined herein; and

with the proviso that the compounds of Formula (III) must contain at least one nitrite, nitrate, thionitrite or thionitrate group.

Another embodiment of the present invention describes compounds of Formula (IV) or pharmaceutically acceptable salts thereof:



IV

wherein

R_{18} and R_{19} at each occurrence are each independently a hydrogen, a lower alkyl group, a halogen, a nitro group, an alkoxy group, $-OD_1$, $-NR_{20}R_{21}$, $-O(O)CR_{20}$, $-O(O)COR_{20}$, $-O(O)CNR_{20}R_{21}$, $-N(R_{20})C(O)R_{21}$, $-N(R_{20})C(O)NR_{20}R_{21}$, or $-N(R_{20})C(O)OR_{21}$; or R_{18} and R_{19} when taken together along with the carbons to which they are attached form a heterocyclic ring or a phenyl ring optionally substituted with up to four substituents selected from a hydrogen, a lower alkyl group, a halogen, a nitro group, an alkoxy group, $-OD_1$, $-NR_{20}R_{21}$, $-O(O)CR_{20}$, $-O(O)COR_{20}$, $-O(O)CNR_{20}R_{21}$, $-N(R_{20})C(O)R_{21}$, $-N(R_{20})C(O)NR_{20}R_{21}$, or $-N(R_{20})C(O)OR_{21}$;

R_{20} and R_{21} at each occurrence are each independently a hydrogen, a lower alkyl group, an aryl group, a lower alkylaryl group, or K;

X_4 is $-C(=R_6)R_{22}$, a heterocyclic ring, $-NR_{20}R_{21}$, a halogen, an alkoxy group, an arylalkoxy group, a cycloalkoxy group, a heterocyclicalkoxy group, an


alkylsulfonyl group, an alkylsulfinyl group, an arylsulfonyl group, an arylsulfinyl group, an arylalkylsulfonyl group, an arylalkylsulfinyl group, a heterocyclicsulfonyl group, or a heterocyclicsulfinyl group;

R_{22} is a hydrogen, an alkyl group, an alkoxy group, an aryl group, an alkylaryl group, a heterocyclic ring, an -O-heterocyclic ring, or an alkylheterocyclic ring;

D_1 , R_6 , and K are defined as herein; and

with the proviso that the compounds of Formula (IV) must contain at least one nitrite, nitrate, thionitrite or thionitrate group.




Another embodiment of the present invention describes compounds of Formula (V) or pharmaceutically acceptable salt thereof:



(V)

wherein

X_5 is:



wherein

R_{23} is a hydrogen, a dialkylamino group, $-NR_7R_7'$, or a heterocyclic ring;

R_{24} is a hydrogen or halogen;

R_{25} is a hydrogen, $-OD_1$, or lower alkyl- OD_1 ;

R_{27} at each occurrence is independently a hydrogen or an alkoxy group;

5 R_{28} , R_{29} , and R_{30} are each independently a hydrogen, a lower alkyl group, a dialkylamino group, a heterocyclic ring, or a lower alkyl- OD_1 ;

R_{31} is a hydrogen, a dialkylamino group, or an alkoxy group;

R_{33} is a hydrogen or a lower alkyl group;


n is an integer from 0 to 1;

10 R_7 , R_7' , D_1 and q are as defined herein; and

with the proviso that the compounds of Formula (V) must contain at least one nitrite, nitrate, thionitrite or thionitrate group.

Another embodiment of the present invention describes compounds of Formula (VI) or pharmaceutically acceptable salts thereof:

15



VI

wherein


20 A_4 , A_5 , and A_6 are each independently a sulfur or CR_{34} with the proviso that one of A_4 , A_5 , or A_6 is a sulfur and the other two are CR_{34} ;

R_{34} at each occurrence is independently a hydrogen, a halogen, a cyano, a nitro, a trifluoromethyl, a lower alkyl group, a heterocyclic ring, a lower alkyl- OD_1 , an alkoxy, a haloalkoxy, an alkylthio, an alkylsulfinyl, an alkylsulfonyl, an alkylcarbonyl, an alkoxy carbonyl, a carbamoyl, a N-alkylcarbamoyl, a N,N-di-
25 alkylcarbamoyl, an ester, a cycloalkyl, an aryl, an alkylaryl, an aryloxy, an arylalkoxyoxy, an arylamino, an alkylaryl amino, an arylthio, an arylsulfonyl, an arylsulfinyl, or a sulfonamido;

R_{35} and R_{36} are each independently a hydrogen or a lower alkyl group; or R_{35} and R_{41} taken together with the carbon chain to which they are attached form

a cycloalkyl ring;

R_{26} is:



wherein

5 X_6 is nitrogen, and Y_6 is CR_{37} ; or X_6 is CR_{37} , and Y_6 is nitrogen;

R_{37} is a hydrogen, a halogen, a lower alkyl group, a trifluoromethyl, an alkoxy group, a haloalkoxy group, an aryl group, an arylalkoxy group, a heterocyclic ring, or an aryloxy;

Z_6 is $-NR_{38}R_{39}$, SR_{40} , or an arylalkoxy group;

10 R_{38} and R_{39} are each independently a hydrogen, a lower alkyl group, an aryl group, an alkylaryl group, or a cycloalkyl group; or R_{38} and R_{39} taken together with the nitrogen to which they are attached form a heterocyclic ring;

R_{40} is a hydrogen, a halogen, a lower alkyl group, an alkylaryl group, an alkenyl group, or a haloalkyl group;

15 R_{41} , R_{42} , R_{43} , and R_{44} are each independently a hydrogen, a halogen, a lower alkyl group, an alkoxy group, a haloalkoxy group, an alkoxyaryl group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a cyano group, $-Y-OD_1$, $Y-SD_1$, $-Y-NR_{20}R_{21}$, $-Y-O(O)CR_{20}$, $-Y-O(O)CNR_{20}R_{21}$, $-Y-N(R_{20})C(O)R_{21}$, or $-Y-N(R_{20})S(O)_2R_{21}$;


20 Y is $-(CH_2)_a-$ or a phenyl group;

R_{45} and R_{46} are each independently a hydrogen, a lower alkyl group, a cycloalkyl group, an alkenyl group, or an alkynyl group;

D_1 , R_{20} , R_{21} , and a are as defined herein; and

25 with the proviso that the compounds of Formula (VI) must contain at least one nitrite, nitrate, thionitrite or thionitrate group.

Another embodiment of the present invention describes compounds of Formula (VII) or pharmaceutically acceptable salts thereof:



VII


5 wherein

R_{60} is a lower alkyl group, an aryl group, a haloalkyl group, a lower alkyl- OD_1 , or heterocyclic ring;


A_7 is oxygen or $-ND_1$;

X_7 is a hydrogen or a halogen;

10 Y_7 is:



or X_7 , A_7 , and Y_7 taken together along with the carbons to which they are attached is:



15 wherein

R_{61} is a hydrogen, a halogen, a lower alkyl group, $-OD_1$, or $-NHC(O)O$ -lower alkyl;

R_{62} is a hydrogen, a halogen, or a lower alkyl group; and

D_1 is as defined herein.

5 Compounds of the present invention that have one or more asymmetric carbon atoms may exist as the optically pure enantiomers, pure diastereomers, mixtures of enantiomers, mixtures of diastereomers, racemic mixtures of enantiomers, diastereomeric racemates or mixtures of diastereomeric racemates. It is to be understood that the present invention anticipates and includes within
10 its scope all such isomers and mixtures thereof.

It is also to be understood that the present invention is intended to include within its scope compounds which may exist in more than one resonance form and the effects that the resonance form may have on the positions at D_1 substituents designated in the structures described herein.

15 Another embodiment of the present invention provides processes for making the novel compounds of the present invention and to the intermediates useful in such processes. The reactions are performed in solvents appropriate to the reagents and materials used are suitable for the transformations being effected. It is understood by one skilled in the art of organic synthesis that the
20 functionality present in the molecule must be consistent with the chemical transformation proposed. This will, on occasion, necessitate judgment by the routineer as to the order of synthetic steps, protecting groups required, and deprotection conditions. Substituents on the starting materials may be incompatible with some of the reaction conditions required in some of the
25 methods described, but alternative methods and substituents compatible with the reaction conditions will be readily apparent to skilled practitioners in the art. The use of sulfur, oxygen and nitrogen protecting groups is well known in the art for protecting thiol, alcohol, and amino groups against undesirable reactions during a synthetic procedure and many such protecting groups are known and
30 described by, for example, Greene and Wuts, *Protective Groups in Organic Synthesis*, Third Edition, John Wiley & Sons, New York (1999).

The chemical reactions described herein are generally disclosed in terms of their broadest application to the preparation of the compounds of this invention.

Occasionally, the reactions may not be applicable as described to each compound included within the disclosed scope. The compounds for which this occurs will be readily recognized by one skilled in the art. In all such cases, either the reactions can be successfully performed by conventional modifications known to one skilled in the art, *e.g.*, by appropriate protection of interfering groups, by changing to alternative conventional reagents, by routine modification of reaction conditions, and the like, or other reactions disclosed herein or otherwise conventional, will be applicable to the preparation of the corresponding compounds of this invention. In all preparative methods, all starting materials are known or readily prepared from known starting materials.

The compounds of Formulas (I), (II), (III), (IV), (V), (VI) and (VII) can be synthesized by one skilled in the art following the methods and examples described herein. The synthesis of the parent proton pump inhibitors (i.e. non-nitrosated and non-nitrosylated proton pump inhibitors) are disclosed in, for example, U. S. Patent Nos. 4,045,564, 4,255,431, 4,634,710, 4,758,579, 4,839,365, 4,873,337, 4,981,861, 5,149,702, 5,554,631, 5,703,097 and 5,945,425 and in EP 0 045 200 A1, EP 0 221 041 A2, EP 0 246 774 A1 and EP 0 254 588 A1 and in WO 98/54172 for the compounds of Formula (I); and in U. S. Patent Nos. 4,806,549, 4,806,550 and 5,952,504 and in EP 0 259 174 A1 and in WO 89/08104 and WO 92/12969 for the compounds of Formula (II); and in U. S. Patent Nos. 5,686,458, 5,750,531 and 5,990,311 and in WO 98/18784 and WO 98/43968 for the compounds of Formula (III); and in U. S. Patent No. 5,677,302 for the compounds of Formula (IV); and in WO 97/32854 for the compounds of Formula (V); and in U. S. Patent Nos. 4,818,760, 4,845,118, 4,871,734, 4,956,366 and 5,114,955 and in EP 0 234 485 A1 for the compounds of Formula (VI); and in U. S. Patent Nos. 5,439,917 and 5,665,730 and in EP 0 033 094 B1 and in WO 95/27714 for the compounds of Formula (VII); the disclosures of each of which are incorporated by reference herein in their entirety.

The nitrosated and nitrosylated proton pump inhibitors of the present invention can be synthesized as shown in reaction Schemes 1 through 7 presented herein. The parent proton pump inhibitor compounds can be nitrosated and/or nitrosylated through one or more sites such as oxygen, sulfur, carbon and/or nitrogen using the methods described in the examples herein and

using conventional methods known to one skilled in the art. For example, known methods for nitrosating and nitrosylating compounds are described in U.S. Patent Nos. 5,380,758 and 5,703,073; WO 97/27749; WO 98/19672; and Oae et al, *Org. Prep. Proc. Int.*, 15(3):165-198 (1983), the disclosures of each of which are
5 incorporated by reference herein in their entirety. The methods of nitrosating and/or nitrosylating the compounds described in the examples herein and in these references can be applied by one skilled in the art to produce any of the nitrosated and/or nitrosylated proton pump inhibitors described herein.


Nitroso or nitro compounds of formula (I), where X is a -ONO, -SNO, or
10 -ONO₂ group and R₁, R₂, R₃, R₄, R₅, R₁₀, R₁₁, A, B, J, R_e, R_f, and p are as defined herein, and a nitrite, nitrate, or thionitrite containing carbamate is representative of the D₁ group, as defined herein, may be prepared as shown in Scheme 1. The substituted imidazole nitrogen group of formula 1 is converted to the anion by treatment with one equivalent of a strong non-nucleophilic base, such as sodium
15 hydride or potassium hydride, in an aprotic solvent, such as tetrahydrofuran (THF) or dimethylformamide (DMF). The carbamate of formula IA, IB, or IC where p, X, R_e and R_f are as defined herein, is prepared by reacting the imidazole anion with a suitably functionalized chloroformate in an inert solvent, such as THF or DMF. Typically the coupling reaction is performed at a temperature
20 ranging from -78 °C to room temperature. Preferred methods for the formation of chloroformates are reacting one equivalent of a X functionalized alcohol with one equivalent of phosgene at a temperature ranging from -78 °C to 0 °C in an inert solvent, such as ether or THF and in the presence of an amine base, such as pyridine or triethylamine. Removal of the amine hydrochloride by filtration
25 affords a solution of the desired chloroformate which may be used directly or concentrated and redissolved in the anhydrous solution of choice prior to the coupling reaction with the imidazole anion to afford the carbamate of formula IA, IB, or IC.

On occasion it might be desirable to nitrosylate the alcohol or thiol after
30 coupling a chloroformate to the imidazole anion. The chloroformate would be prepared by reacting phosgene with an alcohol containing a protected alcohol or thiol moiety. Preferred protecting groups for an alcohol moiety are silyl ethers, such as a trimethylsilyl ether, a tert-butyldimethylsilyl ether, or a tert-

butyldiphenylsilyl ether. After formation of the carbamate, deprotection of the hydroxyl moiety (fluoride ion is the preferred method for removing silyl ether protecting groups) followed by reaction with a suitable nitrosylating agent, such as thionyl chloride nitrite, thionyl dinitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as dichloromethane, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine affords the compound of formula IA. Preferred protecting groups for the thiol moiety are as a thioester, such as a thioacetate or a thiobenzoate or as a disulfide. Deprotection of the thiol moiety (zinc in dilute aqueous acid, triphenylphosphine in water and sodium borohydride are preferred methods for reducing disulfide groups while aqueous base or sodium methoxide in methanol is typically used to hydrolyze thioesters) followed by reaction with a suitable nitrosylating agent such, as thionyl chloride nitrite, thionyl dinitrite, a lower alkyl nitrite, such as tert-butyl nitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as methylene chloride, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula IB. Nitrosation of the carbamate product may be accomplished by first converting the deprotected alcohol to a leaving group such as a mesylate or a tosylate. This reaction is typically performed at a temperature of 0 °C to room temperature in an inert solvent, such as ether, THF, or dichloromethane with the alcohol, methansulfonyl chloride or para-toluensulfonyl chloride, and an amine base, such as triethylamine or pyridine, as the reactants. The corresponding iodide is then prepared by reacting the mesylate or tosylate with sodium iodide in acetone. The halide may also be formed from the alcohol by treatment of the hydroxyl moiety with a phosphorus reagent, such as triphenylphosphine, in the presence of a halide source, such as carbon tetrabromide or N-iodosuccimide, in an inert solvent, such as THF. Treatment of the bromide or iodide with silver nitrate in an inert solvent, such as acetonitrile, affords the compound of formula IC. Alternatively, the halide containing carbamate may be formed directly by preparing a halide containing chloroformate from a halide containing alcohol. Preferred halides are bromide and iodide. Coupling of the imidazole anion with the halide containing

chloroformate followed by reaction of the carbamate product with silver nitrate in an inert solvent, such as acetonitrile, affords the compound of formula IC.

Scheme 1



Nitroso or nitro compounds of formula (II), where X is a -ONO, -SNO, or -ONO₂ group and R₈, R₉, R₇₁, D₁, J, R_e, R_f, o, p, and q are as defined herein, and a nitrite, nitrate, or thionitrite containing alkoxyethyl ester is representative of the

10 K group, as defined herein, may be prepared as shown in Scheme 2. The hydroxyl group of formula 2 is converted to the ester of formula IIA, IIB, or IIC, where p, R_e, R_f and X are as defined herein, by reaction with an appropriate

15 nitrite, thionitrite, or nitrate containing activated acylating agent. Preferred methods for the formation of esters are reacting the alcohol with the preformed acid chloride or anhydride of the nitrite, thionitrite, or nitrate containing acid.

Preferred methods for preparing acid chlorides are treating the carboxylic acid with oxalyl chloride and a catalytic amount of DMF in an inert solvent, such as ether, THF, dichloromethane, or toluene. Preferred methods for preparing

20 mixed anhydride are reacting the carboxylic acid with a chloroformate such as isobutylchloroformate in the presence of an amine base, such as triethylamine in an inert solvent, such as ether, THF, dichloromethane, or toluene.


Alternatively, the alcohol and nitrite, thionitrite, or nitrate containing acid may be condensed in the presence of a dehydrating agent, such as dicyclohexylcarbodiimide (DCC) or 1-ethyl-3 (3-dimethylaminopropyl)

25 carbodiimide hydrochloride (EDAC·HCl) with or without a catalyst, such as 4-dimethylaminopyridine (DMAP) or 1-hydroxybenzotriazole (HOBt).

On occasion it might be desirable to nitrosylate the alcohol or thiol after coupling the activated acylating agent to the alcohol. The activated acylating agent would be prepared from an acid containing a protected alcohol or thiol moiety. Preferred protecting groups for an alcohol moiety are silyl ethers, such as a trimethylsilyl ether, a tert-butyldimethylsilyl ether, or a tert-butyldiphenylsilyl ether. After formation of the ester, deprotection of the hydroxyl moiety (fluoride ion is the preferred method for removing silyl ether protecting groups) followed by reaction with a suitable nitrosylating agent, such as thionyl chloride nitrite, thionyl dinitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as dichloromethane, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula IIA. Preferred protecting groups for the thiol moiety are as a thioester, such as a thioacetate or a thiobenzoate, or as a disulfide. Deprotection of the thiol moiety (zinc in dilute aqueous acid, triphenylphosphine in water and sodium borohydride are preferred methods for reducing disulfide groups while aqueous base or sodium methoxide in methanol is typically used to hydrolyze thioesters) followed by reaction with a suitable nitrosylating agent such, as thionyl chloride nitrite or thionyl dinitrite, a lower alkyl nitrite, such as tert-butyl nitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as methylene chloride, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula IIB. Nitrosation of the ester product may be accomplished by first converting the deprotected alcohol to a leaving group, such as a mesylate or a tosylate. This reaction is typically performed at a temperature of 0 °C to room temperature in an inert solvent, such as ether, THF, or dichloromethane with the alcohol, methanesulfonyl chloride or para-toluensulfonyl chloride, and an amine base, such as triethylamine or pyridine, as the reactants. The corresponding iodide is then prepared by reacting the mesylate or tosylate with sodium iodide in acetone. The halide may also be formed from the alcohol by treatment of the hydroxyl moiety with a phosphorus reagent, such as triphenylphosphine, in the presence of a halide source, such as carbon tetrabromide or N-iodosuccinimide, in an inert solvent, such as THF. Treatment of the bromide or iodide with silver nitrate in an inert solvent, such as acetonitrile affords the compound of formula IIC.

Alternatively, the halide containing ester may be formed directly by preparing a halide containing active acylating agent from a halide containing acid. Preferred halides are bromide and iodide. Coupling of the alcohol with the halide containing active acylating agent followed by reaction of the ester product with silver nitrate in an inert solvent, such as acetonitrile affords the compound of formula IIC. Preferred coupling methods for the formation of esters from alcohols are those methods described herein (e.g. with the preformed acid chloride or anhydride or with the carboxylic acid and a dehydration agent, such as DCC or EDAC·HCl).

Scheme 2



Nitroso or nitro compounds of formula (III), where X is a -ONO, -SNO, or -ONO₂ group and R₁₄, R₁₇, A₁, A₂, A₃, R_e, R_f, Y₃, and p, are as defined herein, and a nitrite, nitrate, or thionitrite containing acyloxymethyl ester is representative of the R₁₃ group may be prepared as shown in Scheme 3. The hydroxyl group of formula 3 is converted to the ester of formula IIIA, IIB, or IIC, where p, R_e, R_f and X are as defined herein, by reaction with an appropriate nitrite, thionitrite, or nitrate containing activated acylating agent. Preferred methods for the formation of esters are reacting the alcohol with the preformed acid chloride or symmetrical anhydride of the nitrite, thionitrite, or nitrate containing acid. Preferred methods for preparing acid chlorides are treating the carboxylic acid with oxalyl chloride and a catalytic amount of DMF in an inert solvent, such as ether, THF,


dichloromethane, or toluene. Preferred methods for preparing mixed anhydride are reacting the carboxylic acid with a chloroformate, such as isobutylchloroformate, in the presence of an amine base, such as triethylamine in an inert solvent solvent, such as ether, THF, dichloromethane, or toluene.

5 Alternatively, the alcohol and nitrite, thionitrite, or nitrate containing acid may be condensed in the presence of a dehydrating agent, such as DCC or EDAC·HCl with or without a catalyst, such as DMAP or HOBt.

On occasion it might be desirable to nitrosylate the alcohol or thiol after coupling the activated acylating agent to the alcohol. The activated acylating
10 agent would be prepared from an acid containing a protected alcohol or thiol moiety. Preferred protecting groups for an alcohol moiety are silyl ethers, such as a trimethylsilyl ether, a tert-butyldimethylsilyl ether, or a tert-butyldiphenylsilyl ether. After formation of the ester, deprotection of the hydroxyl moiety (fluoride ion is the preferred method for removing silyl ether protecting groups) followed
15 by reaction with a suitable nitrosylating agent, such as thionyl chloride nitrite, thionyl dinitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as dichloromethane, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine affords the compound of formula IIIA. Preferred protecting groups for the thiol moiety are as a thioester, such as a
20 thioacetate or a thiobenzoate or as a disulfide. Deprotection of the thiol moiety (zinc in dilute aqueous acid, triphenylphosphine in water and sodium borohydride are preferred methods for reducing disulfide groups while aqueous base or sodium methoxide in methanol is typically used to hydrolyze thioesters) followed by reaction with a suitable nitrosylating agent such, as thionyl chloride
25 nitrite or thionyl dinitrite, a lower alkyl nitrite, such as tert-butyl nitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as methylene chloride, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula IIIB. Nitrosation of the ester product may be accomplished by first converting the
30 deprotected alcohol to a leaving group, such as a mesylate or a tosylate. This reaction is typically performed at a temperature of 0 °C to room temperature in an inert solvent, such as ether, THF, or dichloromethane with the alcohol, methansulfonyl chloride or para-toluensulfonyl chloride, and an amine base,

such as triethylamine or pyridine as the reactants. The corresponding iodide is then prepared by reacting the mesylate or tosylate with sodium iodide in acetone. The halide may also be formed from the alcohol by treatment of the hydroxyl moiety with a phosphorus reagent, such as triphenylphosphine, in the presence of a halide source, such as carbon tetrabromide or N-iodosuccinimide, in an inert solvent, such as THF. Treatment of the bromide or iodide with silver nitrate in an inert solvent, such as acetonitrile, affords the compound of formula IIIC. Alternatively, the halide containing ester may be formed directly by preparing a halide containing active acylating agent from a halide containing acid. Preferred halides are bromide and iodide. Coupling of the alcohol with the halide containing active acylating agent followed by reaction of the ester product with silver nitrate in an inert solvent, such as acetonitrile, affords the compound of formula IIIC. Preferred coupling methods for the formation of esters from alcohols are those methods described herein (e.g. with the preformed acid chloride or anhydride or with the carboxylic acid and a dehydration agent, such as DCC or EDAC·HCl).

Scheme 3



20

Nitroso or nitro compounds of formula (IV), where X is a -ONO, -SNO, or

-ONO₂ group and R₁₈, R₁₉, R_e, R_f, and p, are as defined herein, and a nitrite, nitrate, or thionitrite containing carboxylic acid ester is representative of the X₄ group may be prepared as shown in Scheme 4. The acid of the formula 4 is converted to the ester of formula IVA, IVB, or IVC, where p, R_e, R_f and X are as


5 defined herein, by reaction with an appropriate nitrite, thionitrite, or nitrate containing alcohol. Preferred methods for the preparation of esters are initially forming the mixed anhydride via reaction of 4 with a chloroformate, such as isobutylchloroformate, in the presence of a non nucleophilic base, such as triethylamine, in an anhydrous inert solvent, such as dichloromethane, diethylether or THF. The mixed anhydride is then reacted with the nitrite, 10 thionitrite, or nitrate containing alcohol preferably in the presence of a condensation catalyst, such as DMAP. Alternatively, the acid 4 may be first converted to the acid chloride by treatment with oxalyl chloride in the presence of a catalytic amount of DMF. The acid chloride is then reacted with the nitrite, 15 thionitrite, or nitrate containing alcohol preferably in the presence of a condensation catalyst, such as DMAP, and a tertiary amine base, such as triethylamine, to afford the ester of formula IVA, IVB, or IVC. Alternatively, the acid 4 and nitrite, thionitrite, or nitrate containing alcohol may be coupled to afford the ester of formula IVA, IVB, or IVC by treatment with a dehydration 20 agent, such as DCC or EDAC·HCl, with or without a catalyst, such as DMAP or HOBt.

On occasion it might be desirable to nitrosylate the alcohol or thiol after coupling the acid to the alcohol. The ester would be prepared by reacting the carboxylic acid with an alcohol containing a protected alcohol or thiol moiety. 25 Preferred protecting groups for an alcohol moiety are silyl ethers, such as a trimethylsilyl ether, a tert-butyldimethylsilyl ether, or a tert-butyldiphenylsilyl ether. After coupling the acid and alcohol moieties, deprotection of the hydroxyl moiety (fluoride ion is the preferred method for removing silyl ether protecting groups) followed by reaction with a suitable nitrosylating agent, such as thionyl 30 chloride nitrite, thionyl dinitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as dichloromethane, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula IVA. Preferred protecting groups for the thiol moiety are as a

thioester, such as a thioacetate or a thiobenzoate or as a disulfide. Deprotection of the thiol moiety (zinc in dilute aqueous acid, triphenylphosphine in water and sodium borohydride are preferred methods for reducing disulfide groups while aqueous base or sodium methoxide in methanol is typically used to hydrolyze thioesters) followed by reaction with a suitable nitrosylating agent such, as thionyl chloride nitrite, thionyl dinitrite, a lower alkyl nitrite, such as tert-butyl nitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as methylene chloride, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula IVB.

Nitrosation of the ester product containing a deprotected alcohol moiety may be accomplished by first converting the alcohol to a leaving group, such as a mesylate or a tosylate. This reaction is typically performed at a temperature of 0 °C to room temperature in an inert solvent, such as ether, THF, or dichloromethane with the alcohol, methanesulfonyl chloride or para-toluensulfonyl chloride, and an amine base, such as triethylamine or pyridine, as the reactants. The corresponding iodide is then prepared by reacting the mesylate or tosylate with sodium iodide in acetone. The halide may also be formed from the alcohol by treatment of the hydroxyl moiety with a phosphorus reagent, such as triphenylphosphine, in the presence of a halide source, such as carbon tetrabromide or N-iodosuccinimide, in an inert solvent, such as THF. Treatment of the bromide or iodide with silver nitrate in an inert solvent, such as acetonitrile, affords the compound of formula IVC. Alternatively, the halide containing ester may be formed directly by preparing a halide containing active acylating agent from a halide containing acid. Preferred halides are bromide and iodide. Coupling of the alcohol with the halide containing active acylating agent followed by reaction of the ester product with silver nitrate affords the compound of formula IVC. Preferred coupling methods for the formation of esters from alcohols are those methods described herein (e.g. with the preformed acid chloride or anhydride or with the carboxylic acid and a dehydration agent, such as DCC or EDAC·HCl).

Scheme 4



Nitroso or nitro compounds of formula (V), where X is a -ONO, -SNO, or -ONO₂ group and R₂₃, R₂₄, R₂₅, R₂₇, R₂₉, R₃₀, R_e, R_f, n and p, are as defined herein, and a nitrite, nitrate, or thionitrite containing ester of a substituted pyridine is representative of the X₅ group may be prepared as shown in Scheme 5. The hydroxyl group of formula 5 is converted to the ester of formula VA, VB, or VC, where p, R_e, R_f and X are as defined herein, by reaction with an appropriate


nitrite, thionitrite, or nitrate containing activated acylating agent. Preferred methods for the formation of esters are reacting the alcohol with the preformed acid chloride or symmetrical anhydride of the nitrite, thionitrite, or nitrate containing acid. Preferred methods for preparing acid chlorides are treating the carboxylic acid with oxalyl chloride and a catalytic amount of DMF in an inert solvent, such as ether, THF, dichloromethane, or toluene. Preferred methods for preparing mixed anhydride are reacting the carboxylic acid with a chloroformate, such as isobutylchloroformate, in the presence of an amine base, such as triethylamine, in an inert solvent solvent, such as ether, THF, dichloromethane, or toluene. Alternatively, the alcohol and nitrite, thionitrite, or nitrate containing acid may be condensed in the presence of a dehydrating agent, such as DCC or EDAC·HCl, with or without a catalyst, such as DMAP or HOBT.

On occasion it might be desirable to nitrosylate the alcohol or thiol after coupling the activated acylating agent to the alcohol. The activated acylating agent would be prepared from an acid containing a protected alcohol or thiol

moiety. Preferred protecting groups for an alcohol moiety are silyl ethers, such as a trimethylsilyl ether, a tert-butyldimethylsilyl ether, or a tert-butyldiphenylsilyl ether. After formation of the ester, deprotection of the hydroxyl moiety (fluoride ion is the preferred method for removing silyl ether protecting groups) followed
5 by reaction with a suitable nitrosylating agent, such as thionyl chloride nitrite, thionyl dinitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as dichloromethane, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine affords the compound of formula VA. Preferred protecting groups for the thiol moiety are as a thioester, such as a
10 thioacetate or a thiobenzoate or as a disulfide. Deprotection of the thiol moiety (zinc in dilute aqueous acid, triphenylphosphine in water and sodium borohydride are preferred methods for reducing disulfide groups while aqueous base or sodium methoxide in methanol is typically used to hydrolyze thioesters) followed by reaction with a suitable nitrosylating agent, such as thionyl chloride
15 nitrite, thionyl dinitrite, a lower alkyl nitrite, such as tert-butyl nitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as methylene chloride, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula VB. Nitrosation of the ester product may be accomplished by first converting the
20 deprotected alcohol to a leaving group, such as a mesylate or a tosylate. This reaction is typically performed at a temperature of 0 °C to room temperature in an inert solvent, such as ether, THF, or dichloromethane with the alcohol, methanesulfonyl chloride or para-toluensulfonyl chloride, and an amine base, such as triethylamine or pyridine as the reactants. The corresponding iodide is
25 then prepared by reacting the mesylate or tosylate with sodium iodide in acetone. The halide may also be formed from the alcohol by treatment of the hydroxyl moiety with a phosphorus reagent, such as triphenylphosphine in the presence of a halide source, such as carbon tetrabromide or N-iodosuccinimide in an inert solvent, such as THF. Treatment of the bromide or iodide with silver nitrate in
30 an inert solvent, such as acetonitrile, affords the compound of formula VC. Alternatively, the halide containing ester may be formed directly by preparing a halide containing active acylating agent from a halide containing acid. Preferred halides are bromide and iodide. Coupling of the alcohol with the halide

containing active acylating agent followed by reaction of the ester product with silver nitrate in an inert solvent, such as acetonitrile, affords the compound of formula VC. Preferred coupling methods for the formation of esters from alcohols are those methods described herein (e.g. with the preformed acid chloride or anhydride or with the carboxylic acid and a dehydration agent, such as DCC or EDAC·HCl).

Scheme 5




Nitroso or nitro compounds of formula (VI), where X is a -ONO, -SNO, or -ONO₂ group and R₂₆, R₃₅, R₃₆, A, A₄, A₅, A₆, R_e, R_f, and p are as defined herein, and a nitrite, nitrate, or thionitrite containing carbamate is representative of the D₁ group, as defined herein, may be prepared as shown in Scheme 6. The substituted imidazole nitrogen group of formula 6 is converted to the anion by treatment with one equivalent of a strong non-nucleophilic base, such as sodium hydride or potassium hydride in an aprotic solvent, such as THF or DMF. The carbamate of formula VIA, VIB, or VIC where p, X, R_e and R_f are as defined herein, is prepared by reacting the imidazole anion with a suitably functionalized chloroformate in an inert solvent, such as THF or DMF. Typically the coupling reaction is performed at a temperature ranging between -78 °C and room temperature. Preferred methods for the formation of chloroformates are reacting one equivalent of X functionalized alcohol with one equivalent of phosgene at a temperature ranging from -78 °C to 0 °C in an inert solvent, such as ether or THF,

and in the presence of an amine base, such as pyridine or triethylamine. Removal of the amine hydrochloride by filtration affords a solution of the desired chloroformate which may be used directly or concentrated and redissolved in the anhydrous solution of choice prior to the coupling reaction with the imidazole anion to afford the carbamate of the formula VIA, VIB, or VIC.

On occasion it might be desirable to nitrosylate the alcohol or thiol after coupling a chloroformate to the imidazole anion. The chloroformate would be prepared by reacting phosgene with an alcohol containing a protected alcohol or thiol moiety. Preferred protecting groups for an alcohol moiety are silyl ethers, such as a trimethylsilyl ether, a tert-butyldimethylsilyl ether, or a tert-butyldiphenylsilyl ether. After formation of the carbamate, deprotection of the hydroxyl moiety (fluoride ion is the preferred method for removing silyl ether protecting groups) followed by reaction with a suitable nitrosylating agent, such as thionyl chloride nitrite, thionyl dinitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as dichloromethane, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula VIA. Preferred protecting groups for the thiol moiety are as a thioester, such as a thioacetate or a thiobenzoate or as a disulfide. Deprotection of the thiol moiety (zinc in dilute aqueous acid, triphenylphosphine in water and sodium borohydride are preferred methods for reducing disulfide groups while aqueous base or sodium methoxide in methanol is typically used to hydrolyze thioesters) followed by reaction with a suitable nitrosylating agent, such as thionyl chloride nitrite, thionyl dinitrite, a lower alkyl nitrite, such as tert-butyl nitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as methylene chloride, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula VIB. Nitrosation of the carbamate product may be accomplished by first converting the deprotected alcohol to a leaving group, such as a mesylate or a tosylate. This reaction is typically performed at a temperature of 0 °C to room temperature in an inert solvent, such as ether, THF, or dichloromethane, with the alcohol, methansulfonyl chloride or para-toluensulfonyl chloride, and an amine base, such as triethylamine or pyridine, as

the reactants. The corresponding iodide is then prepared by reacting the mesylate or tosylate with sodium iodide in acetone. The halide may also be formed from the alcohol by treatment of the hydroxyl moiety with a phosphorus reagent, such as triphenylphosphine, in the presence of a halide source, such as carbon tetrabromide or N-iodosuccinimide, in an inert solvent, such as THF. Treatment of the bromide or iodide with silver nitrate in an inert solvent such as acetonitrile affords the compound of formula VIC. Alternatively, the halide containing carbamate may be formed directly by preparing a halide containing chloroformate from a halide containing alcohol. Preferred halides are bromide and iodide. Coupling of the imidazole anion with the halide containing chloroformate followed by reaction of the carbamate product with silver nitrate in an inert solvent such as acetonitrile affords the compound of formula VIC.

Scheme 6



6

VIA X = ONO
 VIB X = SNO
 VIC X = ONO₂


Nitroso or nitro compounds of formula (VII), where X is a -ONO, -SNO, or -ONO₂ group and R₆₀, A₇, X₇, Y₇, R_e, R_f, and p, are as defined herein, and a nitrite, nitrate, or thionitrite containing acyl group is representative of the D group may be prepared as shown in Scheme 7. The hydroxyl group of formula 7 is converted to the ester of formula VIIA, VIIB, or VIIC, where p, R_e, R_f and X are as defined herein, by reaction with an appropriate nitrite, thionitrite, or nitrate containing activated acylating agent. Preferred methods for the formation of esters are reacting the alcohol with the preformed acid chloride or symmetrical

anhydride of the nitrite, thionitrite, or nitrate containing acid. Preferred methods for preparing acid chlorides are treating the carboxylic acid with oxalyl chloride and a catalytic amount of DMF in an inert solvent, such as ether, THF, dichloromethane, or toluene. Preferred methods for preparing mixed anhydride
5 are reacting the carboxylic acid with a chloroformate, such as isobutylchloroformate, in the presence of an amine base, such as triethylamine, in an inert solvent solvent, such as ether, THF, dichloromethane, or toluene. Alternatively, the alcohol and nitrite, thionitrite, or nitrate containing acid may be condensed in the presence of a dehydrating agent, such as DCC or EDAC·HCl
10 with or without a catalyst, such as DMAP or HOBt.

On occasion it might be desirable to nitrosylate the alcohol or thiol after coupling the activated acylating agent to the alcohol. The activated acylating agent would be prepared from an acid containing a protected alcohol or thiol moiety. Preferred protecting groups for an alcohol moiety are silyl ethers, such as
15 a trimethylsilyl ether, a tert-butyldimethylsilyl ether, or a tert-butyldiphenylsilyl ether. After formation of the ester, deprotection of the hydroxyl moiety (fluoride ion is the preferred method for removing silyl ether protecting groups) followed by reaction with a suitable nitrosylating agent, such as thionyl chloride nitrite, thionyl dinitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous
20 solvent, such as dichloromethane, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine affords the compound of formula VIIA. Preferred protecting groups for the thiol moiety are as a thioester, such as a thioacetate or a thiobenzoate or as a disulfide. Deprotection of the thiol moiety (zinc in dilute aqueous acid, triphenylphosphine in water and sodium
25 borohydride are preferred methods for reducing disulfide groups while aqueous base or sodium methoxide in methanol is typically used to hydrolyze thioesters) followed by reaction with a suitable nitrosylating agent, such as thionyl chloride nitrite, thionyl dinitrite, a lower alkyl nitrite, such as tert-butyl nitrite, or nitrosonium tetrafluoroborate, in a suitable anhydrous solvent, such as
30 methylene chloride, THF, DMF, or acetonitrile, with or without an amine base, such as pyridine or triethylamine, affords the compound of formula VIIB. Nitrosation of the ester product may be accomplished by first converting the deprotected alcohol to a leaving group, such as a mesylate or a tosylate. This

reaction is typically performed at a temperature of 0 °C to room temperature in an inert solvent, such as ether, THF, or dichloromethane with the alcohol, methansulfonyl chloride or para-toluensulfonyl chloride, and an amine base, such as triethylamine or pyridine as the reactants. The corresponding iodide is then prepared by reacting the mesylate or tosylate with sodium iodide in acetone. The halide may also be formed from the alcohol by treatment of the hydroxyl moiety with a phosphorus reagent, such as triphenylphosphine in the presence of a halide source, such as carbon tetrabromide or N-iodosuccinimide in an inert solvent, such as THF. Treatment of the bromide or iodide with silver nitrate in an inert solvent, such as acetonitrile, affords the compound of formula VIIC. Alternatively, the halide containing ester may be formed directly by preparing a halide containing active acylating agent from a halide containing acid. Preferred halides are bromide and iodide. Coupling of the alcohol with the halide containing active acylating agent followed by reaction of the ester product with silver nitrate in an inert solvent, such as acetonitrile, affords the compound of formula VIIC. Preferred coupling methods for the formation of esters from alcohols are those methods described herein (e.g. with the preformed acid chloride or anhydride or with the carboxylic acid and a dehydration agent, such as DCC or EDAC·HCl).

Scheme 7



The compounds of the present invention include proton pump inhibitors, such as those described herein, which have been nitrosated and/or nitrosylated through one or more sites such as oxygen (hydroxyl condensation), sulfur (sulfhydryl condensation), carbon and/or nitrogen. The nitrosated and/or

nitrosylated proton pump inhibitors of the present invention are capable of donating, transferring and/or releasing a biologically active form of nitrogen monoxide (i.e., nitric oxide).

5 Nitrogen monoxide can exist in three forms: NO⁻ (nitroxyl), NO[•] (uncharged nitric oxide) and NO⁺ (nitrosonium). NO[•] is a highly reactive short-lived species that is potentially toxic to cells. This is critical because the pharmacological efficacy of NO depends upon the form in which it is delivered. In contrast to the nitric oxide radical (NO[•]), nitrosonium (NO⁺) does not react with O₂ or O₂⁻ species, and functionalities capable of transferring and/or releasing
10 NO⁺ and NO⁻ are also resistant to decomposition in the presence of many redox metals. Consequently, administration of charged NO equivalents (positive and/or negative) is a more effective means of delivering a biologically active NO to the desired site of action.

Compounds contemplated for use in the present invention (e.g., proton
15 pump inhibitors optionally substituted with one or more NO and/or NO₂ groups) can be used in combination with nitric oxide and compounds that release nitric oxide (i.e., compounds that release nitric oxide or otherwise directly or indirectly deliver or transfer nitric oxide to a site of its activity, such as on a cell membrane, *in vivo*, and/or elevate or stimulate production of endogenous nitric
20 oxide or EDRF *in vivo* and/or is a substrate for nitric oxide synthase). "In combination," as used herein can mean that (i) the proton pump inhibitor, optionally substituted with at least one NO and/or NO₂ group, and nitric oxide donor can be present together in the same composition; (ii) the proton pump inhibitor, optionally substituted with at least one NO and/or NO₂ group, and
25 nitric oxide donor can be administered separately; and/or (iii) the proton pump inhibitor, optionally substituted with at least one NO and/or NO₂ group, and nitric oxide donor can be together in the form of a kit.

The term "nitric oxide" encompasses uncharged nitric oxide (NO[•]) and charged nitrogen monoxide species, preferably charged nitrogen monoxide
30 species, such as nitrosonium ion (NO⁺) and nitroxyl ion (NO⁻). The reactive form of nitric oxide can be provided by gaseous nitric oxide. The nitrogen monoxide releasing, delivering or transferring compounds have the structure F-NO, wherein F is a nitrogen monoxide releasing, delivering or transferring

moiety, and include any and all such compounds which provide nitrogen monoxide to its intended site of action in a form active for its intended purpose. The term "NO adducts" encompasses any nitrogen monoxide releasing, delivering or transferring compounds, including, for example, S-nitrosothiols, nitrites, nitrates, S-nitrothiols, sydnonimines, 2-hydroxy-2-nitrosohydrazines (NONOates), (E)-alkyl-2-[(E)-hydroxyimino]-5-nitro-3-hexene amines or amides, nitrosoamines, furoxans as well as substrates for the endogenous enzymes which synthesize nitric oxide. The "NO adducts" can be mono-nitrosylated, poly-nitrosylated, mono-nitrosated and/or poly-nitrosated or a combination thereof at a variety of naturally susceptible or artificially provided binding sites for biologically active forms of nitrogen monoxide.

One group of NO adducts is the S-nitrosothiols, which are compounds that include at least one -S-NO group. These compounds include S-nitroso-polypeptides (the term "polypeptide" includes proteins and polyamino acids that do not possess an ascertained biological function, and derivatives thereof); S-nitrosylated amino acids (including natural and synthetic amino acids and their stereoisomers and racemic mixtures and derivatives thereof); S-nitrosylated sugars; S-nitrosylated, modified and unmodified, oligonucleotides (preferably of at least 5, and more preferably 5-200 nucleotides); straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted S-nitrosylated hydrocarbons; and S-nitroso heterocyclic compounds. S-nitrosothiols and methods for preparing them are described in U.S. Patent Nos. 5,380,758 and 5,703,073; WO 97/27749; WO 98/19672; and Oae et al, *Org. Prep. Proc. Int.*, 15(3):165-198 (1983), the disclosures of each of which are incorporated by reference herein in their entirety.

Another embodiment of the present invention is S-nitroso amino acids where the nitroso group is linked to a sulfur group of a sulfur-containing amino acid or derivative thereof. Such compounds include, for example, S-nitroso-N-acetylcysteine, S-nitroso-captopril, S-nitroso-N-acetylpenicillamine, S-nitroso-homocysteine, S-nitroso-cysteine and S-nitroso-glutathione.

Suitable S-nitrosylated proteins include thiol-containing proteins (where the NO group is attached to one or more sulfur groups on an amino acid or amino acid derivative thereof) from various functional classes including

enzymes, such as tissue-type plasminogen activator (TPA) and cathepsin B; transport proteins, such as lipoproteins; heme proteins, such as hemoglobin and serum albumin; and biologically protective proteins, such as immunoglobulins, antibodies and cytokines. Such nitrosylated proteins are described in WO 93/09806, the disclosure of which is incorporated by reference herein in its entirety. Examples include polynitrosylated albumin where one or more thiol or other nucleophilic centers in the protein are modified.

Other examples of suitable S-nitrosothiols include:

- (i) $\text{HS}(\text{C}(\text{R}_e)(\text{R}_f))_m\text{SNO}$;
 - (ii) $\text{ONS}(\text{C}(\text{R}_e)(\text{R}_f))_m\text{R}_e$; and
 - (iii) $\text{H}_2\text{N}-\text{CH}(\text{CO}_2\text{H})-(\text{CH}_2)_m-\text{C}(\text{O})\text{NH}-\text{CH}(\text{CH}_2\text{SNO})-\text{C}(\text{O})\text{NH}-\text{CH}_2-\text{CO}_2\text{H}$;
- wherein m is an integer from 2 to 20; R_e and R_f are each independently a hydrogen, an alkyl, a cycloalkoxy, a halogen, a hydroxy, an hydroxyalkyl, an alkoxyalkyl, an arylheterocyclic ring, an alkylaryl, a cycloalkylalkyl, a heterocyclicalkyl, an alkoxy, a haloalkoxy, an amino, an alkylamino, a dialkylamino, an arylamino, a diarylamino, an alkylarylamino, an alkoxyhaloalkyl, a haloalkoxy, a sulfonic acid, a sulfonic ester, an alkylsulfonic acid, an arylsulfonic acid, an arylalkoxy, an alkylthio, an arylthio, a cycloalkylthio, a cycloalkenyl, a cyano, an aminoalkyl, an aminoaryl, an aryl, an arylalkyl, an alkylaryl, a carboxamido, an alkylcarboxamido, an arylcarboxamido, an amidyl, a carboxyl, a carbamoyl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarbonyl, an arylcarbonyl, an ester, a carboxylic ester, an alkylcarboxylic ester, an arylcarboxylic ester, a haloalkoxy, a sulfonamido, an alkylsulfonamido, an arylsulfonamido, a sulfonic ester, a urea, a phosphoryl, a nitro, -T-Q, or $(\text{C}(\text{R}_e)(\text{R}_f))_k-\text{T}-\text{Q}$, or R_e and R_f taken together with the carbons to which they are attached form a carbonyl, a methanthial, a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group; Q is -NO or -NO₂; and T is independently a covalent bond, a carbonyl, an oxygen, -S(O)_o- or -N(R_s)R_t-, wherein o is an integer from 0 to 2, R_s is a lone pair of electrons, a hydrogen or an alkyl group; R_t is a hydrogen, an alkyl, an aryl, an alkylcarboxylic acid, an aryl carboxylic acid, an alkylcarboxylic ester, an arylcarboxylic ester, an alkylcarboxamido, an arylcarboxamido, an alkylaryl, an alkylsulfinyl, an alkylsulfonyl, an arylsulfinyl, an arylsulfonyl, a sulfonamido, a carboxamido, a carboxylic ester, an amino alkyl, an amino aryl,

$-\text{CH}_2-\text{C}(\text{T-Q})(\text{R}_e)(\text{R}_f)$, or $-(\text{N}_2\text{O}_2^-) \cdot \text{M}^+$, wherein M^+ is an organic or inorganic cation; with the proviso that when R_f is $-\text{CH}_2-\text{C}(\text{T-Q})(\text{R}_e)(\text{R}_f)$ or $-(\text{N}_2\text{O}_2^-) \cdot \text{M}^+$; then "-T-Q" can be a hydrogen, an alkyl group, an alkoxyalkyl group, an aminoalkyl group, a hydroxy group or an aryl group.

5 In cases where R_e and R_f are a heterocyclic ring or taken together R_e and R_f are a heterocyclic ring, then R_f can be a substituent on any disubstituted nitrogen contained within the radical wherein R_f is as defined herein.

Nitrosothiols can be prepared by various methods of synthesis. In general, the thiol precursor is prepared first, then converted to the S-nitrosothiol derivative by nitrosation of the thiol group with NaNO_2 under acidic conditions (pH is about 2.5) which yields the S-nitroso derivative. Acids which can be used for this purpose include aqueous sulfuric, acetic and hydrochloric acids. The thiol precursor can also be nitrosylated by reaction with an organic nitrite such as tert-butyl nitrite, or a nitrosonium salt such as nitrosonium tetrafluoroborate in an inert solvent.

10 Another group of NO adducts for use in the present invention, where the NO adduct is a compound that donates, transfers or releases nitric oxide, include compounds comprising at least one ON-O-, ON-N- or ON-C- group. The compounds that include at least one ON-O-, ON-N- or ON-C- group are preferably ON-O-, ON-N- or ON-C-polypeptides (the term "polypeptide" includes proteins and polyamino acids that do not possess an ascertained biological function, and derivatives thereof); ON-O-, ON-N- or ON-C-amino acids (including natural and synthetic amino acids and their stereoisomers and racemic mixtures); ON-O-, ON-N- or ON-C-sugars; ON-O-, ON-N- or ON-C-

25 modified or unmodified oligonucleotides (comprising at least 5 nucleotides, preferably 5-200 nucleotides); ON-O-, ON-N- or ON-C- straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted hydrocarbons; and ON-O-, ON-N- or ON-C-heterocyclic compounds.

30 Another group of NO adducts for use in the present invention include nitrates that donate, transfer or release nitric oxide, such as compounds comprising at least one $\text{O}_2\text{N-O-}$, $\text{O}_2\text{N-N-}$, $\text{O}_2\text{N-S-}$ or $\text{O}_2\text{N-C-}$ group. Preferred among these compounds are $\text{O}_2\text{N-O-}$, $\text{O}_2\text{N-N-}$, $\text{O}_2\text{N-S-}$ or $\text{O}_2\text{N-C-}$ polypeptides (the term "polypeptide" includes proteins and also polyamino acids that do not

possess an ascertained biological function, and derivatives thereof); O_2N-O- , O_2N-N- , O_2N-S- or O_2N-C- amino acids (including natural and synthetic amino acids and their stereoisomers and racemic mixtures); O_2N-O- , O_2N-N- , O_2N-S- or O_2N-C- sugars; O_2N-O- , O_2N-N- , O_2N-S- or O_2N-C- modified and unmodified
 5 oligonucleotides (comprising at least 5 nucleotides, preferably 5-200 nucleotides); O_2N-O- , O_2N-N- , O_2N-S- or O_2N-C- straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted hydrocarbons; and O_2N-O- , O_2N-N- , O_2N-S- or O_2N-C- heterocyclic compounds. Preferred examples of compounds comprising at least one O_2N-O- , O_2N-N- , O_2N-S- or
 10 O_2N-C- group include isosorbide dinitrate, isosorbide mononitrate, clonitrate, erythrityltetranitrate, mannitol hexanitrate, nitroglycerin, pentaerythritoltetranitrate, pentrinitrol and propatylnitrate.

Another group of NO adducts are N-oxo-N-nitrosoamines that donate, transfer or release nitric oxide and are represented by the formula:
 15 $R^1R^2-N(O-M^+)-NO$, where R^1 and R^2 are each independently a polypeptide, an amino acid, a sugar, a modified or unmodified oligonucleotide, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted hydrocarbon, or a heterocyclic group, and where M^+ is an organic or inorganic cation, such as, for example, an alkyl substituted ammonium cation
 20 or a Group I metal cation.

Another group of NO adducts are thionitrates that donate, transfer or release nitric oxide and are represented by the formula: $R^1-(S)-NO_2$, where R^1 is a polypeptide, an amino acid, a sugar, a modified or unmodified oligonucleotide, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted
 25 or unsubstituted hydrocarbon, or a heterocyclic group. Preferred are those compounds where R^1 is a polypeptide or hydrocarbon with a pair or pairs of thiols that are sufficiently structurally proximate, i.e., vicinal, that the pair of thiols will be reduced to a disulfide. Compounds which form disulfide species release nitroxyl ion (NO^-) and uncharged nitric oxide (NO^\bullet). Compounds where
 30 the thiol groups are not sufficiently close to form disulfide bridges generally provide nitric oxide as the NO^- form and not as the uncharged NO^\bullet form.

The present invention is also directed to compounds that stimulate endogenous NO or elevate levels of endogenous endothelium-derived relaxing

factor (EDRF) *in vivo* or are substrates for the enzyme, nitric oxide synthase. Such compounds include, for example, L-arginine, L-homoarginine, and N-hydroxy-L-arginine, including their nitrosated and nitrosylated analogs (e.g., nitrosated L-arginine, nitrosylated L-arginine, nitrosated N-hydroxy-L-arginine, nitrosylated N-hydroxy-L-arginine, nitrosated L-homoarginine and nitrosylated L-homoarginine), precursors of L-arginine and/or physiologically acceptable salts thereof, including, for example, citrulline, ornithine, glutamine, lysine, polypeptides comprising at least one of these amino acids, inhibitors of the enzyme arginase (e.g., N-hydroxy-L-arginine and 2(S)-amino-6-boronoheptanoic acid) and the substrates for nitric oxide synthase, cytokines, adenosin, bradykinin, calreticulin, bisacodyl, and phenolphthalein. EDRF is a vascular relaxing factor secreted by the endothelium, and has been identified as nitric oxide (NO) or a closely related derivative thereof (Palmer et al, *Nature*, 327:524-526 (1987); Ignarro et al, *Proc. Natl. Acad. Sci. USA*, 84:9265-9269 (1987)).

Another aspect of the invention provides methods for preventing and/or treating gastrointestinal disorders by administering to the patient in need thereof a therapeutically effective amount of the compounds and/or compositions described herein. Such gastrointestinal disorders include, for example, inflammatory bowel disease, Crohn's disease, irritable bowel syndrome, ulcerative colitis, peptic ulcers, stress ulcers, bleeding peptic ulcers, duodenal ulcers, infectious enteritis, colitis, diverticulitis, gastric hyperacidity, dyspepsia, gastroparesis, Zollinger-Ellison syndrome, gastroesophageal reflux disease, *Helicobacter Pylori* associated disease, short-bowel (anastomosis) syndrome, hypersecretory states associated with systemic mastocytosis or basophilic leukemia and hyperhistaminemia that result, for example, from neurosurgery, head injury, severe body trauma or burns. In one aspect of the invention, the patient can be administered at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention to prevent and/or treat the gastrointestinal disorder. In another aspect of the invention, the patient can be administered at least one antacid and at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention to prevent or treat the gastrointestinal disorder. In another aspect of the invention, the patient can be administered at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention and at

least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to prevent and/or treat the gastrointestinal disorder. In still another aspect of the invention, the patient can be administered at least one antacid, at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention, and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to prevent and/or treat the gastrointestinal disorder. In yet another aspect of the present invention, the patient can be administered at least one proton pump inhibitor and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to prevent and/or treat the gastrointestinal disorder. In yet another aspect of the present invention, the patient can be administered at least one antacid, at least one proton pump inhibitor, and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to prevent and/or treat the gastrointestinal disorder.

The antacid, proton pump inhibitor that is optionally substituted with at least one NO and/or NO₂ group, and/or the nitric oxide donor can be administered separately or as components of the same composition. These compounds and/or compositions can also be provided in the form of a pharmaceutical kit. The proton pump inhibitors substituted with at least one NO and/or NO₂ group and preferred nitric oxide donors are described in detail herein. Appropriate antacids for use in this aspect of the invention include any antacid known in the art, including, for example, aluminum hydroxide, magnesium hydroxide, magnesium carbonate, calcium carbonate and co-dried gels, such as, for example, aluminum hydroxide-magnesium carbonate co-dried gel.

Another aspect of the present invention provides methods to improve the gastroprotective properties, anti-*Helicobacter* properties and antacid properties of proton pump inhibitors by administering to a patient in need thereof a therapeutically effective amount of the compounds and/or compositions described herein. In one aspect of the invention, the patient can be administered

at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention to improve the gastroprotective properties, anti-*Helicobacter* properties and antacid properties of the proton pump inhibitor. In another aspect of the invention, the patient can be administered a bismuth-complex comprising at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention to improve the gastroprotective properties, anti-*Helicobacter* properties and antacid properties of the proton pump inhibitor. In another aspect of the invention, the patient can be administered at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to improve the gastroprotective properties, anti-*Helicobacter* properties and antacid properties of the proton pump inhibitor. In another aspect of the invention, the patient can be administered a bismuth complex comprising at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to improve the gastroprotective properties, anti-*Helicobacter* properties and antacid properties of the proton pump inhibitor. In yet another aspect of the invention, the patient can be administered at least one proton pump inhibitor and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to improve the gastroprotective properties, anti-*Helicobacter* properties and antacid properties of the proton pump inhibitor. In yet another aspect of the present invention, the patient can be administered a bismuth-complex comprising at least one proton pump inhibitor and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to improve the gastroprotective properties, anti-*Helicobacter* properties and antacid properties of the proton pump inhibitor.

The bismuth-containing reagent, proton pump inhibitor, that is optionally, substituted with at least one NO and/or NO₂ group, and/or nitric oxide donor can be administered separately or as components of the same

composition. The proton pump inhibitors, optionally substituted with at least one NO and/or NO₂ group, and nitric oxide donors are described in detail herein. Bismuth complexes are prepared by boiling the aqueous solution of the free base of the proton pump inhibitor with at least one bismuth-containing reagent, including, for example, bismuth citrate, bismuth salicylate, bismuth tartaric acid or mixtures thereof as described in U. S. Patent No. 5,403,830 and in Ivanov et al, *J. Pharm. Pharmacol.*, 48:297-301 (1996), the disclosures of which are incorporated by reference herein in their entirety.

Another aspect of the present invention provides methods to facilitate ulcer healing and decrease the recurrence of ulcers by administering to a patient in need thereof a therapeutically effective amount of the compounds and/or compositions described herein. In one aspect of the invention, the patient can be administered at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention to facilitate ulcer healing and decrease the recurrence of ulcers. In another aspect of the invention, the patient can be administered at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase to facilitate ulcer healing and decrease the recurrence of ulcers. In yet another aspect of the invention, the patient can be administered at least one proton pump inhibitor and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to facilitate ulcer healing and decrease the recurrence of ulcers. The proton pump inhibitor, that is optionally, substituted with at least one NO and/or NO₂ group, and/or nitric oxide donor can be administered separately or as components of the same composition. The proton pump inhibitor, optionally substituted with at least one NO and/or NO₂ group, and nitric oxide donors are described in detail herein.

Another aspect of the present invention provides methods to decrease or reverse gastrointestinal toxicity and facilitate ulcer healing resulting from, for example, the administration of nonsteroidal antiinflammatory drugs (NSAIDs), selective COX-2 inhibitors, and the like. In particular, the present invention provides methods of administering a therapeutically effective amount of the

compounds and/or compositions described herein, and, optionally, administering a therapeutically effective amount of at least one NSAID or selective COX-2 inhibitor. In one aspect of the invention, the patient can be administered at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention, and, optionally, at least one NSAID and/or selective COX-2 inhibitor, to decrease or reverse gastrointestinal toxicity and/or to facilitate ulcer healing resulting from the NSAID and/or selective COX-2 inhibitor treatment. In another aspect of the invention, the patient can be administered at least one NSAID and/or selective COX-2 inhibitor with a therapeutically effective amount of at least one nitrosated and/or nitrosylated proton pump inhibitor of the invention and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to decrease or reverse gastrointestinal toxicity and/or to facilitate ulcer healing resulting from the NSAID and/or selective COX-2 inhibitor treatment. In yet another aspect of the present invention, the patient can be administered at least one NSAID and/or selective COX-2 inhibitor with a therapeutically effective amount of at least one proton pump inhibitor and at least one compound that donates, transfers or releases nitric oxide, or elevates endogenous levels of nitric oxide or EDRF, or is a substrate for nitric oxide synthase, to decrease or reverse gastrointestinal toxicity and/or to facilitate ulcer healing resulting from the NSAID and/or selective COX-2 inhibitor treatment. The NSAID and/or selective COX-2 inhibitor, nitrosated and/or nitrosylated proton pump inhibitor, proton pump inhibitor, and/or nitric oxide donor can be administered separately or as components of the same composition. These compounds and/or compositions can also be provided in the form of a pharmaceutical kit.

The compounds and compositions of the present invention can be used in this aspect of the invention with any NSAID and selective COX-2 inhibitor known in the art. Such NSAIDs include, for example, aspirin (e.g., acetylsalicylic acid), salicylate esters and salts, acetate esters of salicylic acid, difluorophenyl derivatives (e.g., diflunisal), salicylsalicylic acids (e.g., salsalate), salts of salicylic acids (e.g., sodium salicylate), salicylamide, sodium thiosalicylate, choline salicylate, magnesium salicylate, combinations of choline and magnesium

salicylates, 5-aminosalicylic acid (e.g., mesalamine), salicylazosulfapyridine (e.g., sulfasalazine), methylsalicylate, and the like.

Another group of NSAIDs are the pyrazolon derivatives, which include, for example, phenylbutazone, oxyphenbutazone, antipyrine, aminopyrine, dipyrone and apazone (azapropazone). Another group of NSAIDs are the para-aminophenol derivatives, which are the so-called "coal tar" analgesics, including, for example, phenacetin and its active metabolite acetaminophen. Another group of compounds include indomethacin, a methylated indole derivative, and the structurally related compound sulindac. Yet another group of compounds is the fenamates which are derivatives of N-phenylanthranilic acid (e.g., mefenamic, meclofenamic, flufenamic, tolfenamic and etofenamic acids). Another contemplated NSAID is tolmetin.

Another group of NSAIDs are the propionic acid derivatives. Principal members of this group are, for example, ibuprofen, naproxen, flurbiprofen, fenoprofen and ketoprofen. Other members of this group include, for example, fenbufen, pirprofen, oxaprozin, indoprofen and tiaprofenic acid.

Still other NSAIDs are piroxicam, ampiroxicam, oxicam derivatives (which are a class of antiinflammatory enolic acids), tenoxicam, tenidap, diclofenac (one of the series of phenylacetic acid derivatives that have been developed as antiinflammatory agents). Other NSAIDs include etodolac and nabumentone.

Selective COX-2 inhibitors are disclosed in, for example, U. S. Patent Nos. 5,681,842, 5,750,558, 5,756,531, 5,776,984 and in WO 98/39330, WO 99/10331 and WO 99/10332 assigned to Abbott Laboratories; and in WO 98/50075 assigned to Algos Pharmaceutical Corporation; and in U. S. Patent No. 5,980,905 assigned to AMBI Inc.; and in U. S. Patent Nos. 5,776,967, 5,824,699, 5,830,911 and in WO 98/04527 and WO 98/21195 assigned to American Home Products Corporation; and in WO 99/18960 assigned to Astra Pharmaceuticals Ltd.; and in U. S. Patent No. 5,905,089 assigned to Board of Supervisors of Louisiana State University; and in WO 97/13767 assigned to Chemisch Pharmazeutische Forschungsgesellschaft MBH; and in WO 96/10021 assigned to The Du Pont Merck Pharmaceutical Company; and in WO 99/13799 assigned to Euro-Celtique; and in U. S. Patent No. 5,134,142 and in WO 99/15505 assigned to Fujisawa Pharmaceutical Co. Ltd.; and

in U. S. Patent Nos. 5,344,991, 5,393,790, 5,521,207, 5,596,008, 5,616,601, 5,620,999, 5,633,272, 5,643,933, 5,686,470, 5,696,143, 5,700,816, 5,859,257, 5,972,986, 5,990,148 and in WO 94/15932, WO 94/27980, WO 95/15316, WO 96/16934, WO 96/25405, WO 96/38418, WO 96/38442, WO 96/41645, WO 97/38986, WO 98/06708, WO 98/43649, 5 WO 98/47509, WO 98/47890 and WO 99/22720 assigned to G.D. Searle & Co.; and in WO 96/31509 and WO 99/12930 assigned to Glaxo Group Limited; and in WO 97/34882 assigned to Grupo Farmaceutico Almirall; and in WO 97/03953 assigned to Hafslund Nycomed Pharma AG; and in U. S. Patent No. 5,945,539, 5,994,381 and in EP 0 745 596 A1 assigned to Japan Tobacco, Inc.; and in U. S. Patent No. 10 5,686,460, 5,807,873 and in WO 97/37984 and WO 99/21585 assigned to Laboratoires USPA; and in U. S. Patent Nos. 5,585,504, 5,840,924, 5,883,267, 5,925,631 and in WO 97/44027, WO 97/44028, WO 97/45420, WO 98/00416, WO 98/47871, WO 99/15503, WO 99/15513, WO 99/20110, WO 99/45913 and WO 99/55830 assigned to Merck & Co. Inc.; and in U. S. Patent Nos. 5,409,944, 15 5,436,265, 5,474,995, 5,536,752, 5,550,142, 5,510,368, 5,521,213, 5,552,422, 5,604,253, 5,604,260, 5,639,780, 5,677,318, 5,691,374, 5,698,584, 5,710,140, 5,733,909, 5,789,413, 5,817,700, 5,840,746, 5,849,943, 5,861,419, 5,994,379 and in EP 0 788 476 B1, EP 0 863 134 A1 and in WO 94/20480, WO 94/13635, WO 94/26731, WO 95/00501, WO 96/19469, WO 96/37467, WO 97/14691, WO 97/16435, WO 97/28120, WO 97/28121, 20 WO 97/36863, WO 98/03484, WO 98/43966, WO 99/14194, WO 99/14195 and WO 99/23087 assigned to Merck Frosst Canada & Co., and in WO 99/59635 assigned to Merck Sharp & Dohme Limited; and in U. S. Patent No. 5,380,738 assigned to Monsanto Company; and in WO 99/33796 assigned to Nissin Food Products Co. Ltd.; and in U. S. Patent No. 5,783,597 assigned to Ortho Pharmaceutical Corporation; and in WO 98/07714 assigned to Oxis International Inc.; and in EP 0 25 937 722 A1 and in WO 98/50033 and WO 99/05104 assigned to Pfizer Inc.; and in U. S. Patent No. 5,908,858 assigned to Sankyo Company Limited; and in WO 97/25045 assigned to Smithkline Beecham Corporation; and in U. S. Patent No. 5,475,021 assigned to Vanderbilt University; and in WO 99/59634 assigned to 30 Wakamoto Pharmaceutical Co. Ltd., the disclosures of each of which are incorporated by reference herein in their entirety.

Each of the above NSAIDs and selective COX-2 inhibitors is described more fully in the literature, such as in Goodman and Gilman, The

Pharmacological Basis of Therapeutics (8th Edition), McGraw-Hill, pages 617-657 (1993); Merck Index on CD-ROM, Twelfth Edition, Version 12:1, (1996); STN Express, file phar and file registry, the disclosures of which are incorporated by reference herein in their entirety.

5 Other NSAIDs and selective COX-2 inhibitors that can be used in the present invention include those described in U. S. Patent No. 5,703,073, and co-pending Application Serial No. 09/441,891 and 60/171,623, the disclosures of which are incorporated by reference herein in their entirety.

10 Another embodiment of the present invention provides methods to treat *Helicobacter pylori* by administering one or more acid-degradable antibacterial compounds in combination with at least one proton pump inhibitor that is substituted with at least one NO and/or NO₂ group, and, optionally, a therapeutically effective amount of at least one compound that donates, transfers or releases nitric oxide and/or stimulates endogenous production of NO or EDRF
15 *in vivo*. Alternatively, this embodiment provides administering one or more acid-degradable antibacterial compounds in combination with at least one proton pump inhibitor, and, a therapeutically effective amount of at least one compound that donates, transfers or releases nitric oxide and/or stimulates endogenous production of NO or EDRF *in vivo*. U.S. Patent Nos. 5,629,305 and
20 5,599,794, the disclosures of each of which are incorporated by reference herein in their entirety, disclose treating or preventing gastrointestinal disorders resulting from *Helicobacter pylori* by administering proton pump inhibitors in combination with antibacterial compounds. The proton pump inhibitors increase the intra-gastric pH in the stomach, thereby increasing the bioavailability
25 of the acid-labile antibacterial compound. The antibacterial compound(s), proton pump inhibitor(s), optionally substituted with at least one NO and/or NO₂ group, and/or nitric oxide donor(s) can be administered separately, or as components of the same composition. The compounds and/or compositions can also be provided in the form of a kit. The proton pump inhibitors that are
30 optionally substituted with at least one NO group and nitric oxide donors are described herein. The antibacterial compounds contemplated for use in this embodiment of the invention include any acid-degradable antibacterial compound that is known in the art, including, for example antibiotics, such as,

for example, amoxycillin, penicillin, benzylpenicillin, erythromycin base, clarithromycin, and the like.

Yet another embodiment of the present invention provides methods to treat viral infections by administering at least one proton pump inhibitor that is substituted with at least one NO and/or NO₂ group, and, optionally, a therapeutically effective amount of at least one compound that donates, transfers or releases nitric oxide and/or stimulates endogenous production of NO or EDRF *in vivo*. Alternatively, this embodiment provides administering at least one proton pump inhibitor, and, a therapeutically effective amount of at least one compound that donates, transfers or releases nitric oxide and/or stimulates endogenous production of NO or EDRF *in vivo*. U.S. Patent No. 5,945,425, the disclosure of which is incorporated by reference herein in its entirety, discloses treating viral infections by administering proton pump inhibitors. The proton pump inhibitor(s), optionally substituted with at least one NO and/or NO₂ group, and/or nitric oxide donor(s) can be administered separately, or as components of the same composition. The compounds and/or compositions can also be provided in the form of a kit. The proton pump inhibitors that are optionally substituted with at least one NO group and nitric oxide donors are described herein.

The compounds and compositions of the present invention can be administered by any available and effective delivery system including, but not limited to, orally, buccally, parenterally, by inhalation spray, by topical application, by injection, transdermally, or rectally (e.g., by the use of suppositories) in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles, as desired. Parenteral includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques.

Transdermal compound administration, which is known to one skilled in the art, involves the delivery of pharmaceutical compounds via percutaneous passage of the compound into the systemic circulation of the patient. Topical administration can also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. Other components can be incorporated into the transdermal patches as well. For example, compositions

and/or transdermal patches can be formulated with one or more preservatives or bacteriostatic agents including, but not limited to, methyl hydroxybenzoate, propyl hydroxybenzoate, chlorocresol, benzalkonium chloride, and the like.

Dosage forms for topical administration of the compounds and compositions can include creams, sprays, lotions, gels, ointments, eye drops, nose drops, ear drops, and the like. In such dosage forms, the compositions of the invention can be mixed to form white, smooth, homogeneous, opaque cream or lotion with, for example, benzyl alcohol 1% or 2% (wt/wt) as a preservative, emulsifying wax, glycerin, isopropyl palmitate, lactic acid, purified water and sorbitol solution. In addition, the compositions can contain polyethylene glycol 400. They can be mixed to form ointments with, for example, benzyl alcohol 2% (wt/wt) as preservative, white petrolatum, emulsifying wax, and tenox II (butylated hydroxyanisole, propyl gallate, citric acid, propylene glycol). Woven pads or rolls of bandaging material, e.g., gauze, can be impregnated with the compositions in solution, lotion, cream, ointment or other such form can also be used for topical application. The compositions can also be applied topically using a transdermal system, such as one of an acrylic-based polymer adhesive with a resinous crosslinking agent impregnated with the composition and laminated to an impermeable backing.

Solid dosage forms for oral administration can include capsules, tablets, effervescent tablets, chewable tablets, pills, powders, sachets, granules and gels. In such solid dosage forms, the active compounds can be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms can also comprise, as in normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, effervescent tablets, and pills, the dosage forms can also comprise buffering agents. Soft gelatin capsules can be prepared to contain a mixture of the active compounds or compositions of the present invention and vegetable oil. Hard gelatin capsules can contain granules of the active compound in combination with a solid, pulverulent carrier such as lactose, saccharose, sorbitol, mannitol, potato starch, corn starch, amylopectin, cellulose derivatives of gelatin. Tablets and pills can be prepared with enteric coatings.

Liquid dosage forms for oral administration can include pharmaceutically

acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions can also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

5 Suppositories for vaginal or rectal administration of the compounds and compositions of the invention can be prepared by mixing the compounds or compositions with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at room temperature but liquid at body temperature, such that they will melt and release the drug.

10 Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated according to the known art using suitable dispersing agents, wetting agents and/or suspending agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in
15 1,3-butanediol. Among the acceptable vehicles and solvents that can be used are water, Ringer's solution, and isotonic sodium chloride solution. Sterile fixed oils are also conventionally used as a solvent or suspending medium.

 The compositions of this invention can further include conventional excipients, i.e., pharmaceutically acceptable organic or inorganic carrier
20 substances suitable for parenteral application which do not deleteriously react with the active compounds. Suitable pharmaceutically acceptable carriers include, for example, water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, surfactants, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides,
25 petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, and the like. The pharmaceutical preparations can be sterilized and if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavoring and/or aromatic substances and the like which do not deleteriously react with
30 the active compounds. For parenteral application, particularly suitable vehicles consist of solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants. Aqueous suspensions may contain substances which increase the viscosity of the suspension and include, for example, sodium

carboxymethyl cellulose, sorbitol and/or dextran. Optionally, the suspension may also contain stabilizers.

5 The composition, if desired, can also contain minor amounts of wetting agents, emulsifying agents and/or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulations can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like.

10 Various delivery systems are known and can be used to administer the compounds or compositions of the present invention, including, for example, encapsulation in liposomes, microbubbles, emulsions, microparticles, microcapsules and the like. The required dosage can be administered as a single unit or in a sustained release form

15 The bioavailability of the compositions can be enhanced by micronization of the formulations using conventional techniques such as grinding, milling, spray drying and the like in the presence of suitable excipients or agents such as phospholipids or surfactants.

20 The compounds and compositions of the present invention can be formulated as pharmaceutically acceptable salts. Pharmaceutically acceptable salts include, for example, alkali metal salts and addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids include, but are not limited to, hydrochloric, hydrobromic, hydroiodic, nitrous (nitrite salt), nitric (nitrate salt), carbonic, sulfuric, phosphoric acid, and the like. Appropriate organic acids include, but are not limited to, aliphatic, cycloaliphatic, aromatic, heterocyclic, carboxylic and sulfonic classes of organic acids, such as, for example, formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, p-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic,

ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, stearic, algenic, β -hydroxybutyric, cyclohexylaminosulfonic, galactaric and galacturonic acid and the like. Suitable pharmaceutically-acceptable base addition salts include, but are not limited to, 5 metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from primary, secondary and tertiary amines, cyclic amines, N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine and the like. All of these salts may be prepared by conventional means 10 from the corresponding compound by reacting, for example, the appropriate acid or base with the compound.

"Therapeutically effective amount" refers to the amount of the proton pump inhibitor, that is optionally substituted with at least one NO and/or NO₂ group, nitric oxide donor, antacid, bismuth-complex, NSAID, selective COX-2 15 inhibitor and/or acid-degradable antibacterial compound, that is effective to achieve its intended purpose. While individual patient needs may vary, determination of optimal ranges for effective amounts of each of the compounds and compositions is within the skill of the art. Generally, the dosage required to provide an effective amount of the composition, and which can be adjusted by 20 one of ordinary skill in the art will vary, depending on the age, health, physical condition, sex, weight, extent of the dysfunction of the recipient, frequency of treatment and the nature and scope of the dysfunction or disease.

The amount of a given proton pump inhibitor, that is optionally substituted with at least one NO and/or NO₂ group, which will be effective in the 25 treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques, including reference to Goodman and Gilman, *supra*; The Physician's Desk Reference, Medical Economics Company, Inc., Oradell, N.J., 1995; and Drug Facts and Comparisons, Inc., St. Louis, MO, 1993. The precise dose to be used in the 30 formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided by the physician and the patient's circumstances.

The amount of nitric oxide donor in a pharmaceutical composition can be

in amounts of about 0.1 to about 10 times the molar equivalent of the proton pump inhibitor. The usual daily doses of proton pump inhibitors are about 10 mg to about 400 mg per day and the doses of nitric oxide donors in the pharmaceutical composition can be in amounts of about 1 to about 500 mg/kg of body weight daily, preferably about 1 to about 50 mg/kg of body weight daily. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems and are in the same ranges or less than as described for the commercially available compounds in the Physician's Desk Reference, *supra*.

The present invention also provides pharmaceutical kits comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compounds and/or compositions of the present invention, including, at least, one or more of the proton pump inhibitors, that are optionally substituted with at least one NO moiety, and one or more of the NO donors described herein. Such kits can also include, for example, other compounds and/or compositions (e.g., NSAIDs, antacids and/or antibacterial compounds), a device(s) for administering the compounds and/or compositions, and written instructions in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which instructions can also reflect approval by the agency of manufacture, use or sale for human administration.

EXAMPLES

Example 1: {2-[(2-Pyridylmethyl)sulfinyl]benzimidazolyl)methyl-3-[N-[2-methyl-2-(nitrosothio)-propyl]-N-benzylcarbamoyl]propanoate

1a. 2-Methyl-1-(benzylamino)propane-2-thiol

A solution of α,α' -dithiodiisobutyraldehyde (10.31 g, 0.05 mol) and benzylamine (10.71 g, 0.10 mol) in chloroform (150 ml) was heated to reflux for 2 hours. The solution was cooled to room temperature, the chloroform was removed under vacuum and the residue was taken in up methanol (100 ml).

Under ice cooling, sodium borohydride (6 g, 0.16 mol) was then added in portions over 1 hour and the resulting solution was stirred at room temperature for 1 hour. The reaction was concentrated under vacuum and the residue was partitioned between water (300 ml) and ethyl acetate (50 ml). The aqueous phase

was extracted with ethyl ether (2 x 50 ml) and the combined organic phases were washed with brine and dried (Na_2SO_4). The volatiles were evaporated to give an oil (18.45 g, 95%) which was used directly in the next step. ^1H NMR (300 MHz, CDCl_3): δ 1.29 (s, 12 H), 2.57 (s, 4H), 3.81 (s, 4H), 7.21-7.33 (m, 10 H). To a solution of the above oil (13.26 g, 0.034 mol) in ether (70 ml) and liquid ammonia (100 ml) cooled over a dry ice/acetone bath was added sodium until a blue color persisted for 40 min (~2 g of sodium was consumed). Ammonium chloride (15 g) was added and the reaction mixture was left overnight at room temperature. Water (100 ml) was added and the ether layer was separated. The aqueous phase was extracted with ether (2 x 50 ml) and the combined ether layers were washed with brine and dried (Na_2SO_4). The solvent was evaporated to give the title compound (12.83 g, 96.3%) as a colorless oil. ^1H NMR (300 MHz, CDCl_3): δ 1.37 (s, 6 H), 2.61 (s, 2 H), 3.86 (s, 2 H), 7.22-7.36 (m, 5 H); ^{13}C NMR (75 MHz, CDCl_3) δ 30.58, 45.43, 54.24, 62.78, 126.86, 127.94, 128.34, 140.64; LCMS (m/e): 196 (M+1).

1b. 3-[N-(2-Methyl-2-sulfanylpropyl)-N-benzylcarbamoyl]propanoic acid

To a solution of the product of example 1a (4.52 g, 23.14 mmol) in dichloromethane (70 ml) at 0 °C was added succinic anhydride (2.20 g, 21.98 mmol). The reaction was stirred at room temperature overnight, washed with water (100 ml), the aqueous phase was extracted with dichloromethane (2 x 50 ml) and the combined organic phases were washed with brine and dried (Na_2SO_4). The solvent was evaporated under vacuum and the residue triturated with ether/hexane to give the title compound (6.46 g, 99.5%) as a white solid. mp. 123-126°C; ^1H NMR (300 MHz, CDCl_3): δ 1.40 and 1.47 (2 s, 6 H), 1.82 (s, 1H), 2.65-2.86 (m, 4 H), 3.47 and 3.61 (2s, 2 H), 4.89 and 4.93 (2s, 2 H), 7.13-7.54 (m, 5 H), 10.41 (br s, 1 H); ^{13}C NMR (75 MHz, CDCl_3): δ 28.21, 29.39, 31.27, 46.43, 53.01, 58.51, 125.86, 127.50, 128.96, 136.52, 173.49, 177.63; LCMS (m/e): 296 (M+1).

1c. 3-[N-[2-Methyl-2-(nitrosothio)propyl]-N-benzylcarbamoyl]propanoic acid

To a solution of the product of example 1b (4.46 g, 15.1 mmol) in dichloromethane (100 ml) at 0 °C was added dropwise t-butyl nitrite (1.94 ml, 16.6 mmol). The resulting green solution was left at room temperature for 1 hour. The solvent was evaporated and hexane (20 ml) was added to the residue. The mixture was stored at -20 °C for 1 hour, then the hexane was decanted and the solid dried under vacuum to give pure the title compound (4.46 g, 91.1%) as

green crystals. mp. 95-98°C; ¹H NMR (300 MHz, CDCl₃): δ 1.91 and 1.98 (2s, 6 H), 2.63-2.78 (m, 4 H), 4.07 and 4.19 (2 s, 2 H), 4.60 and 4.80 (2 s, 2 H), 7.06-7.37 (m, 5 H); ¹³C NMR (75 MHz, CDCl₃): δ 27.65, 28.26, 29.33, 53.04, 55.74, 58.50, 125.96, 127.75, 129.07, 136.06, 173.81, 177.64; LCMS (m/e): 325 (M+1).

- 5 1d. {2-[(2-Pyridylmethyl)sulfinyl]benzimidazolyl}methyl-3-[N-[2-methyl-2-(nitrosothio)-propyl]-N-benzylcarbamoyl]propanoate

A mixture of timoprazole (0.51 g, 1.98 mmol) and 37% formaldehyde (0.80 ml) in acetonitrile (5 ml) was heated at 70 °C for 15 min. The reaction was evaporated to give a solid that was washed twice with ethyl ether/ethyl acetate (4:1) and dissolved in dichloromethane (10 ml). To this solution was added
 10 Example 1c (0.65 g, 2 mol), 4-(N-N-dimethylamino)pyridine (10 mg) and dicyclohexylcarbodiimide (1M in dichloromethane, 4 ml, 4 mmol). After stirring at room temperature for 3 hours, the reaction was concentrated under vacuum and the residue was purified by flash column chromatography (ethyl
 15 acetate/hexane, 4:1 to 7:1) to afford the title compound (0.65 g, 54.7%) as a green foam. ¹H NMR (300 MHz, CDCl₃): δ 1.82 and 1.91 (2 s, 6 H), 2.63 and 2.75 (2br s, 4 H), 4.08 and 4.13 (2 s, 2 H), 4.70 and 4.85 (2 s, 2H), 4.88 (d, J = 14 Hz, 1 H), 4.92 (d, J = 14 Hz, 1 H), 6.36 (d, J = 11.4 Hz, 1H), 6.42 (d, J = 11.4 Hz, 1H), 7.00 and 7.06 (2 d, J = 6.8 Hz, 2 H), 7.20-7.64 (m, 9 H), 7.81 (d, J = 7.7 Hz, 1H), 8.54 (d, J = 4 Hz, 1 H); ¹³C
 20 NMR (75 MHz, CDCl₃): δ 27.29, 27.94, 28.88, 52.58, 55.29, 58.18, 62.22, 65.04, 110.73, 120.73, 122.97, 123.85, 125.21, 125.30, 125.58, 127.41, 128.76, 135.03, 135.79, 136.66, 141.70, 149.71, 150.20, 151.66, 171.80, 172.76; LCMS (m/e): 594 (M+1).

Example 2: {2-([3-methyl-4-(2,2,2-trifluoroethyl)-2-pyridyl]methyl)sulfinyl}benzimidazolyl-methyl-3-[N-[2-methyl-2-(nitrosothio)-propyl]-N-benzylcarbamoyl]propanoate

- 25 2a {2-([3-methyl-4-(2,2,2-trifluoroethyl)-2-pyridyl]methyl)sulfinyl}benzimidazolyl -methyl-3-[N-[2-methyl-2-(nitrosothio)-propyl]-N-benzylcarbamoyl]propanoate

The procedure described in Example 1d was repeated using lansoprazole (0.739 g, 2 mmol) instead of timoprazole. The title compound (0.82 g, 59.4%) was
 30 obtained as a green foam. ¹H NMR (300 MHz, CDCl₃): δ 1.84 and 1.92 (2 s, 6 H), 2.28 (s, 3 H), 2.65 and 2.76 (2 br s, 4 H), 4.01 and 4.10 (2s, 2 H), 4.39 (q, J = 7.3 Hz, 2 H), 4.56 and 4.72 (2 s, 2 H), 4.96 (d, J = 14 Hz, 1 H), 5.03 (d, J = 14 Hz, 1 H), 6.46 (d, J =

11 Hz, 1 H), 6.54 (d, $J = 11$ Hz, 1 H), 6.65 (d, $J = 5$ Hz, 1 H), 7.00-7.40 (m, 7 H), 7.64 (d, $J = 7.4$ Hz, 1 H), 7.80 (d, $J = 7.5$ Hz, 1 H), 8.25 (d, $J = 5$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3): δ 24.85, 27.47, 28.13, 29.11, 33.82, 52.80, 55.53, 58.37, 59.81, 65.26 (q, $J = 36$ Hz), 65.36, 105.81, 110.85, 120.93, 122.97, 123.92, 125.28, 125.78, 127.58, 128.93, 135.21, 136.01, 141.97, 148.09, 151.30, 152.81, 161.67, 171.96, 172.95; LCMS (m/e): 690 (M+1).

5 **Example 3: 2-[2-(Nitrosothio)adamantan-2-yl]ethyl 2-([3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridyl]methyl)sulfinyl)benzimidazolecarboxylate**

3a. Adamantane-2-thione

Adamantan-2-one (48.46 g, 322.6 mmol) in pyridine (300 mL) was heated to 90 °C and phosphorous pentasulfide (17.84 g, 40.13 mmol) was added. The reaction was maintained at 90 °C for two hours and at room temperature overnight during which time a precipitate formed. The pyridine solution was decanted and concentrated to dryness. The residual semisolid was treated with hexane (400 mL) to give an orange solution with a light brown suspension. The suspension was removed by filtration. The filtrate was concentrated to dryness and dried to vacuum to give an orange solid (50.36 g). This crude product was purified by filtration through a pad of silica gel (hexane). ^1H NMR (CDCl_3 , 300 MHz): δ 3.43 (s, 2H), 2.1-1.9 (m, 12H); ^{13}C NMR (75 MHz, CDCl_3): δ 222.4, 57.5, 41.1, 36.5, 27.4.

20 3b. *tert*-Butyl 2-(2-sulfanyladamantan-2-yl)acetate

To *t*-butyl acetate (25 mL, 21.6 g, 186 mmol) in dry THF (400 mL) at -78 °C was added lithium diisopropylamide monotetrahydrofuran (1.5 M solution in cyclohexane, 100 mL, 150 mmol) under nitrogen and the reaction mixture was stirred at -78 °C for 40 minutes. The product of Example 3a (21.88 g, 131.57 mmol) in THF (400 mL) was added. The cold bath was removed and the reaction was stirred at room temperature for two hours. The reaction was diluted with methylene chloride and 2 M HCl (75 mL) was added. The organic phase was separated, washed with brine (4 x 40 mL), dried (MgSO_4), filtered, and concentrated. The crude product was purified by filtration through a pad of silica gel (5% EtOAc/95% hexane) to give the title compound (34.67 g, 122.7 mmol, 93%). $\text{Rf} = 0.48$ (EtOAc/ hexane 1:19); ^1H NMR (CDCl_3 , 300 MHz): δ 2.87 (s, 2H), 2.47 (d, $J = 11.5$, 2H), 2.38 (s, 1H), 2.11 (d, $J = 11.9$, 2H), 1.98 (s, 2H), 1.96 (m, 2H), 1.84-1.62 (m, 6H), 1.47 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): δ 170.8, 80.7, 54.1, 47.3,

39.0, 38.2, 37.2, 36.6, 34.0, 33.3, 28.2, 27.5, 26.9. APIMS (IS, NH_4OAc) m/e 283 (MH^+); Anal. Calcd for $\text{C}_{16}\text{H}_{26}\text{O}_2\text{S}$ (282.44): C, 68.04; H, 9.28 Found: C, 68.14; H, 9.30.

3c. 2-(2-Sulfanyladamantan-2-yl)ethan-1-ol

To a 0 °C cooled solution of Example 3b (4.1 g, 24.1 mmol) in anhydrous dichloromethane (40 mL) lithium aluminum hydride (1 M solution in THF) (40 mL) was added dropwise over a period of 20 minutes. The reaction mixture was stirred at 0 °C for further 15 minutes and then at room temperature for 30 minutes. The excess LiAlH_4 was destroyed by the addition of ethyl acetate. The reaction mixture was then poured over ice cold water, acidified with 1 N HCl and extracted with dichloromethane (2 x 200 mL). The combined extracts were washed with brine (1 x 75 mL), dried over sodium sulfate, filtered and solvent evaporated at reduced pressure to give the title compound (3.1 g), mp 68-70 °C; ^1H NMR (CDCl_3): δ 1.16-1.9 (m, 11 H), 2.1 (m, 2 H), 2.22 (t, $J = 6.9$ Hz, 3 H), 2.43 (m, 2 H), 3.93 (t, $J = 6.9$ Hz, 2 H); ^{13}C NMR (CDCl_3): δ 26.8, 27.7, 33.2, 33.9, 38.2, 39.1, 43.4, 55.8, 59.4; LRMS (APIMS) (m/z) 230 ($M + 18$) ($M + \text{NH}_4$).

3d. 2-[2-(Nitrosothio)adamantan-2-yl]ethan-1-ol

To a 0 °C cooled solution of Example 3c (1.06 g, 5 mmol) in anhydrous dichloromethane (40 mL) was added t-butyl nitrite (7.5 mmol, 890 μL). The reaction mixture was stirred at 0 °C for 30 minutes and then at room temperature for 30 minutes. The solvent was removed at reduced pressure and product was recrystallized from ethyl ether/hexane to give 1.2 g (80% yield) of the title compound as a green crystalline solid, mp 77-79 °C; ^1H NMR (CDCl_3): δ 1.7-1.74 (m, 2 H), 1.83-1.93 (m, 5 H), 2.06 (m, 3 H), 2.42-2.53 (m, 4 H), 2.99 (t, $J = 7.3$ Hz, 2 H), 3.83 (t, $J = 7.6$ Hz, 2 H); ^{13}C NMR (CDCl_3): δ 27.3, 27.4, 33.2, 33.9, 35.6, 38.97, 39.96, 59.1, 68.2; LRMS (APIMS) (m/z) 259 ($M + 18$) ($M + \text{NH}_4$).

3e. 2-[2-(nitrosothio)adamanta-2-yl]ethyl chlorooate

An ice-cooled solution of 20% phosgene in toluene (2.5 ml, 5.4 mmol) was added dropwise to a solution of Example 3d (0.41 g, 1.7 mmol) and pyridine (0.138 ml, 1.7 mmol) in dichloromethane (4 ml). After stirring at 0 °C for 20 min, the reaction was concentrated using a rotary evaporator. To the residue was added dry ether (20 ml). The reaction mixture was filtered, washed with ether (2 x 5 ml) and the organic layer was concentrated under vacuum to gave the crude product, which was used immediately for next step.

3f. 2-[2-(Nitrosothio)adamantan-2-yl]ethyl 2-([3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridyl]methyl)sulfinyl)benzimidazolecarboxylate

Lansoprazole (0.50 g, 1.35 mmol) was added to a suspension of sodium hydride (39 mg, 1.63 mmol) in N,N-dimethylformamide (3 ml) at room temperature. The resulting clear solution was cooled in an ice bath. A solution of Example 3e in N,N-dimethylformamide (3 ml) was then added. After stirring at 0 °C for 30 min, the reaction mixture was diluted with ethyl ether/ethyl acetate (3:2 v/v) (70 ml) and washed with water (2 x 30 ml) and then brine (20 ml). The combined organic extracts were dried over Na₂SO₄ and evaporated under vacuum. The residue was purified by flash chromatography (SiO₂, 4/1 to 3/1 dichloromethane /acetone) to give the title compound as a green solid (0.52 g, 60.3%). Mp. 125-130°C (dec.)(at 116°C it started to shrink). ¹H NMR (300 MHz, CDCl₃) δ 8.06 (d, J = 5.6 Hz, 1 H), 7.90 (d, J = 7.5 Hz, 1H), 7.82 (d, J = 7.7 Hz, 1 H), 7.48-7.38 (m, 2 H), 6.56 (d, J = 5.6 Hz, 1 H), 4.78 (d, J = 13 Hz, 1H), 4.75 (t, J = 7.8 Hz, 2H), 4.67 (d, J = 13 Hz, 1H), 4.37 (q, J = 7.8 Hz, 2H), 3.37 (t, J = 7.8 Hz, 2H), 2.60-2.46 (m, 4 H), 2.35 (s, 3H), 2.17-1.76 (m, 10 H) ; ¹³C NMR (75 MHz, CDCl₃) δ 161.74, 156.92, 151.19, 149.65, 147.94, 142.77, 133.54, 126.50, 125.24, 124.55, 121.38, 114.61, 105.66, 67.04, 65.88, 65.36 (q, J = 36 Hz), 59.45, 38.74, 35.68, 35.58, 35.52, 33.73, 33.10, 33.08, 27.26, 27.11, 11.37; MS (m/e): 637 (M+1).

Example 4: Comparative *In Vivo* Gastric Lesion Activity

The ethanol/HCl mixture-induced gastric lesion test in rats described by Takeuchi et al, *J. Pharmacol. Exp. Ther.*, 286: 115-121 (1998), was used to evaluate gastric lesion activity. Male Sprague Dawley rats (Charles River Laboratories, Wilmington, MA) weighing 230-250 g were used for the experiments. The rats were housed with laboratory chow and water *ad libitum* prior to the study. The rats were fasted for 24 hours with free access to water and then dosed by oral gavage with vehicle or with the test compounds given at a volume of 0.5 ml/100 g body weight. Thirty minutes after oral dosing all the rats received 1 ml of a solution of 60% ethanol in 150 mM HCl intragastrically. Food was withheld after dosing. Sixty minutes after ethanol/HCl, rats were euthanized by pre-charged CO₂. The stomachs were dissected along the greater curvature, washed with a directed stream of 0.9% saline and pinned open on a sylgard based petri dish for examination of the hemorrhagic lesions. Gastric lesion score was expressed in

mm and calculated by summing the length of each lesion as described by Al-Ghamdi et al, *J. Int. Med. Res.*, 19: 2242 (1991). Results are expressed as the mean \pm standard error of the mean. Statistical analysis were conducted using ANOVA test followed by a Student-Newman-Keuls post-hoc test using the Abacus

5 Concepts, Super Anova computer program (Abacus Concepts, Inc., Berkeley, CA).

Fig. 1 compares the gastric lesion activity of vehicle alone, lansoprazole in vehicle and Example 2 (nitrosylated lansoprazole) in vehicle. Ethanol/HCl mixture produced gastric lesion in the control rats treated with vehicle (0.5% Methocel). Lansoprazole at a dose of 200 μ mol/kg failed to significantly inhibit
10 the formation of gastric lesions. The nitrosylated lansoprazole derivative of Example 2, at 200 μ mol/kg, significantly inhibited the formation of gastric lesions produced by the ethanol/HCl mixture.

The disclosure of each patent, patent application and publication cited or described in the present specification is hereby incorporated by reference herein
15 in its entirety.

Although the invention has been set forth in detail, one skilled in the art will appreciate that numerous changes and modifications can be made to the invention, and that such changes and modifications can be made without departing from the spirit and scope of the present invention.


CLAIMS

What is claimed is:

1. A proton pump inhibitor compound comprising at least one NO group, at least one NO₂ group, or at least one NO and NO₂ group, or a pharmaceutically acceptable salt thereof.

2. The proton pump inhibitor compound of claim 1, wherein the compound comprising at least one NO group, at least one NO₂ group, or at least one NO and NO₂ group is a compound of formula (I), formula (II), formula (III), formula (IV), formula (V), formula (VI) or formula (VII), or a pharmaceutically acceptable salt thereof:

wherein the compound of formula (I) is:



(I)

wherein

A is S, S(O), or S(O)₂;

B is -CNR₇R₇' or nitrogen;

J is CH or nitrogen;


R₁ is a hydrogen, an alkoxy group, a lower alkyl group, or an alkylthio group;

R₂ is a hydrogen, an alkoxy group, a lower alkyl group, an alkylthio group, a haloalkoxy group, an alkoxyalkyl group, -NR₇R₇', -OD₁, or -SD₁; or R₂ and R₁ taken together with the carbon chain to which they are attached form a cycloalkyl ring or a heterocyclic ring; or R₂ and R₃ taken together with the carbon chain to which they are attached form a cycloalkyl ring or a heterocyclic ring;

R₃ and R₁₁ are each independently a hydrogen, an alkoxy group, a lower alkyl group, or an alkylthio group; or R₃ and R₁₁ taken together with the carbon

chain to which they are attached form a cycloalkyl ring or a heterocyclic ring;

R_4 and R_5 are each independently a hydrogen, an alkyl group, a halo group, an alkoxy group, a haloalkyl group, a haloalkoxy group, a cyano group, an aryl group, a heterocyclic ring, $-NR_7R_7'$, $-OD_1$, or $-CO_2R_{12}$; or R_4 and R_5 taken together are:



wherein

R_6 is oxygen or $N=O-R_7$;

R_7 and R_7' are each independently hydrogen, a lower alkyl group or D; or R_7 and R_7' taken together with the nitrogen to which they are attached form a heterocyclic ring;

R_{10} is a hydrogen; or R_{10} and R_1 taken together with the carbon chain to which they are attached form a cycloalkyl ring;

R_{12} is a lower alkyl group or D;

D_1 is a hydrogen or D;

D is Q or K;

Q is $-NO$ or $-NO_2$;

K is $-W_a-E_b-(C(R_e)(R_f))_p-E_c-(C(R_e)(R_f))_x-W_d-(C(R_e)(R_f))_y-W_i-E_j-W_g-(C(R_e)(R_f))_z-$
T-Q;

a, b, c, d, g, i and j are each independently an integer from 0 to 3;

p, x, y and z are each independently an integer from 0 to 10;

W at each occurrence is independently $-C(O)-$, $-C(S)-$, $-T-$, $-(C(R_e)(R_f))_h-$, an alkyl group, an aryl group, a heterocyclic ring, an arylheterocyclic ring, or $-(CH_2CH_2O)_q-$;

E at each occurrence is independently $-T-$, an alkyl group, an aryl group, $-(C(R_e)(R_f))_h-$, a heterocyclic ring, an arylheterocyclic ring, or $-(CH_2CH_2O)_q-$;

h is an integer from 1 to 10;

q is an integer from 1 to 5;

R_e and R_f are each independently a hydrogen, an alkyl, a cycloalkoxy, a halogen, a hydroxy, an hydroxyalkyl, an alkoxyalkyl, an arylheterocyclic ring, an alkylaryl, a cycloalkylalkyl, a heterocyclicalkyl, an alkoxy, a haloalkoxy, an amino, an alkylamino, a dialkylamino, an arylamino, a diarylamino, an alkylaryl amino, an alkoxyhaloalkyl, a haloalkoxy, a sulfonic acid, a sulfonic ester, an alkylsulfonic acid, an arylsulfonic acid, an arylalkoxy, an alkylthio, an arylthio, a cycloalkylthio, a cycloalkenyl, a cyano, an aminoalkyl, an aminoaryl, an aryl, an arylalkyl, an alkylaryl, a carboxamido, an alkylcarboxamido, an arylcarboxamido, an amidyl, a carboxyl, a carbamoyl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarbonyl, an arylcarbonyl, an ester, a carboxylic ester, an alkylcarboxylic ester, an arylcarboxylic ester, a haloalkoxy, a sulfonamido, an alkylsulfonamido, an arylsulfonamido, a sulfonic ester, a urea, a phosphoryl, a nitro, -T-Q, or $(C(R_e)(R_f))_k$ -T-Q, or R_e and R_f taken together with the carbons to which they are attached form a carbonyl, a methanthial, a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group;

k is an integer from 1 to 3;

T at each occurrence is independently a covalent bond, a carbonyl, an oxygen,

-S(O)_o- or -N(R_a)R_i-;


o is an integer from 0 to 2;

R_a is a lone pair of electrons, a hydrogen or an alkyl group;

R_i is a hydrogen, an alkyl, an aryl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarboxylic ester, an arylcarboxylic ester, an alkylcarboxamido, an arylcarboxamido, an alkylaryl, an alkylsulfinyl, an alkylsulfonyl, an arylsulfinyl, an arylsulfonyl, a sulfonamido, a carboxamido, a carboxylic ester, an amino alkyl, an amino aryl, -CH₂-C(T-Q)(R_e)(R_f), or -(N₂O₂)⁻•M⁺, wherein M⁺ is an organic or inorganic cation; with the proviso that when R_i is -CH₂-C(T-Q)(R_e)(R_f) or -(N₂O₂)⁻•M⁺, or R_e or R_f are T-Q or $(C(R_e)(R_f))_k$ -T-Q, then the "-T-Q" subgroup can be a hydrogen, an alkyl, an alkoxy, an alkoxyalkyl, an aminoalkyl, a hydroxy, a heterocyclic ring or an aryl group; and

with the proviso that the compound of formula (I) must contain at least one nitrite, nitrate, thionitrite or thionitrate group;

wherein the compound of formula (II) is:



II

wherein

5 R_8 is a lower alkyl group, an alkoxyalkyl group, an alkylaryl group, a cycloalkyl group, a cycloalkylalkyl group, an aryl group, an alkylaryl group, or K;

R_9 at each occurrence is independently a hydrogen, a lower alkyl group, an alkylthio group, a halogen, a cyano group, an alkanoyl group, a haloalkyl group, a carbamoyl group, $-NR_7D_1$, $-OD_1$, or $-CO_2R_{12}$;


10 R_{71} is a hydrogen, a lower alkyl group, an alkoxy group, or $-OD_1$;

J, K, D_1 , R_7 , R_{12} , q and o are as defined herein; and

with the proviso that the compound of formula (II) must contain at least one nitrite, nitrate, thionitrite or thionitrate group;

wherein the compound of formula (III) is:

15 wherein




III

R_{13} and R_{14} are each independently a hydrogen a lower alkyl group, an alkoxyalkyl, or a lower alkyl- OD_1 ; or R_{13} and R_{14} taken together along with the carbons to which they are attached form a cycloalkyl group or an aryl group;

R_{17} is a hydrogen or a lower alkyl group;

Y_3 is:



wherein

R_{15} is a hydrogen or a lower alkyl group;

R_{16} is a hydrogen, a halogen, or a lower alkyl group;

R_{63} is a lower alkyl group or a phenyl group;

A_1 , A_2 and A_3 comprise the other subunits of a 5- or 6-membered monocyclic aromatic ring and A_1 , A_2 and A_3 are each independently:

(i) CR_o , wherein R_o at each occurrence is hydrogen or $-OD_1$;

(ii) $N-R_p$, wherein R_p at each occurrence is independently a covalent bond to an adjacent ring atom in order to render the ring aromatic, a hydrogen, or K;

(iii) a sulfur atom;


(iv) an oxygen atom; or

(v) $B_a = B_b$, wherein B_a and B_b are each independently a nitrogen atom or CR_o ; wherein R_o at each occurrence is hydrogen or $-OD_1$;

D_1 and K are as defined herein; and

with the proviso that the compound of formula (III) must contain at least one nitrite, nitrate, thionitrite or thionitrate group;

wherein the compound of formula (IV) is:



IV

wherein

5 R_{18} and R_{19} at each occurrence are each independently a hydrogen, a lower alkyl group, a halogen, a nitro group, an alkoxy group, $-OD_1$, $-NR_{20}R_{21}$, $-O(O)CR_{20}$, $-O(O)COR_{20}$, $-O(O)CNR_{20}R_{21}$, $-N(R_{20})C(O)R_{21}$, $-N(R_{20})C(O)NR_{20}R_{21}$, or $-N(R_{20})C(O)OR_{21}$; or R_{18} and R_{19} when taken together along with the carbons to which they are attached form a heterocyclic ring or a phenyl ring optionally
 10 substituted with up to four substituents selected from a hydrogen, a lower alkyl group, a halogen, a nitro group, an alkoxy group, $-OD_1$, $-NR_{20}R_{21}$, $-O(O)CR_{20}$, $-O(O)COR_{20}$, $-O(O)CNR_{20}R_{21}$, $-N(R_{20})C(O)R_{21}$, $-N(R_{20})C(O)NR_{20}R_{21}$, or $-N(R_{20})C(O)OR_{21}$;

R_{20} and R_{21} at each occurrence are each independently a hydrogen, a lower
 15 alkyl group, an aryl group, a lower alkylaryl group, or K;


X_4 is $-C(=R_6)R_{22}$, a heterocyclic ring, $-NR_{20}R_{21}$, a halogen, an alkoxy group, an arylalkoxy group, a cycloalkoxy group, a heterocyclicalkoxy group, an alkylsulfonyl group, an alkylsulfinyl group, an arylsulfonyl group, an arylsulfinyl group, an arylalkylsulfonyl group, an arylalkylsulfinyl group, a
 20 heterocyclicsulfonyl group, or a heterocyclicsulfinyl group;

R_{22} is a hydrogen, an alkyl group, an alkoxy group, an aryl group, an alkylaryl group, a heterocyclic ring, an -O-heterocyclic ring, or an alkylheterocyclic ring;

D_1 , R_6 , and K are defined as herein; and

25 with the proviso that the compound of formula (IV) must contain at least one nitrite, nitrate, thionitrite or thionitrate group;

wherein the compound of formula (V) is:






(V)

wherein

X_5 is:

5



R_{23} is a hydrogen, a dialkylamino group, $-NR_7R_7'$, or a heterocyclic ring;

R_{24} is a hydrogen or halogen;

R_{25} is a hydrogen, $-OD_1$, or lower alkyl- OD_1 ;

10 R_{27} at each occurrence is independently a hydrogen or an alkoxy group;

R_{28} , R_{29} , and R_{30} are each independently a hydrogen, a lower alkyl group, a dialkylamino group, a heterocyclic ring, or a lower alkyl- OD_1 ;

R_{31} is a hydrogen, a dialkylamino group, or an alkoxy group;

R_{33} is a hydrogen or a lower alkyl group;


15 n is an integer from 0 to 1;

R_7 , R_7' , D_1 and q are as defined herein; and

with the proviso that the compound of formula (V) must contain at least one nitrite, nitrate, thionitrite or thionitrate group;

wherein the compound of formula (VI) is:

20



VI


wherein

A_4 , A_5 , and A_6 are each independently a sulfur or CR_{34} with the proviso
5 that one of A_4 , A_5 , or A_6 is a sulfur and the other two are CR_{34} ;

R_{34} at each occurrence is independently a hydrogen, a halogen, a cyano, a
nitro, a trifluoromethyl, a lower alkyl group, a heterocyclic ring, a lower alkyl-
OD₁, an alkoxy, a haloalkoxy, an alkylthio, an alkylsulfinyl, an alkylsulfonyl, an
alkylcarbonyl, an alkoxycarbonyl, a carbamoyl, a N-alkylcarbamoyl, a N,N-di-
10 alkylcarbamoyl, an ester, a cycloalkyl, an aryl, an alkylaryl, an aryloxy, an
arylalkoxyoxy, an arylamino, a alkylarylamino, an arylthio, an arylsulfonyl, an
arylsulfinyl, or a sulfonamido;

R_{35} and R_{36} are each independently a hydrogen or a lower alkyl group; or
 R_{35} and R_{41} taken together with the carbon chain to which they are attached form
15 a cycloalkyl ring;

R_{26} is:



wherein

X_6 is nitrogen, and Y_6 is CR_{37} ; or X_6 is CR_{37} , and Y_6 is nitrogen;

R_{37} is a hydrogen, a halogen, a lower alkyl group, a trifluoromethyl, an
20 alkoxy group, a haloalkoxy group, an aryl group, an arylalkoxy group, a
heterocyclic ring, or an aryloxy;

Z_6 is $-NR_{38}R_{39}$, SR_{40} , or an arylalkoxy group;

R_{38} and R_{39} are each independently a hydrogen, a lower alkyl group, an aryl group, an alkylaryl group, or a cycloalkyl group; or R_{38} and R_{39} taken together with the nitrogen to which they are attached form a heterocyclic ring;

R_{40} is a hydrogen, a halogen, a lower alkyl group, an alkylaryl group, an alkenyl group, or a haloalkyl group;

R_{41} , R_{42} , R_{43} , and R_{44} are each independently a hydrogen, a halogen, a lower alkyl group, an alkoxy group, a haloalkoxy group, an alkoxyaryl group, an alkylthio group, an alkylsulfinyl group, an alkylsulfonyl group, a cyano group, $-Y-OD_1$, $-Y-SD_1$, $-Y-NR_{20}R_{21}$, $-Y-O(O)CR_{20}$, $-Y-O(O)CNR_{20}R_{21}$, $-Y-N(R_{20})C(O)R_{21}$, or $-Y-N(R_{20})S(O)_2R_{21}$;


Y is $-(CH_2)_a-$ or a phenyl group;

R_{45} and R_{46} are each independently a hydrogen, a lower alkyl group, a cycloalkyl group, an alkenyl group, or an alkynyl group;

D_1 , R_{20} , R_{21} , and a are as defined herein; and

with the proviso that the compound of formula (VI) must contain at least one nitrite, nitrate, thionitrite or thionitrate group;

wherein the compound of formula (VII) is:



VII

20

wherein


R_{60} is a lower alkyl group, an aryl group, a haloalkyl group, a lower alkyl- OD_1 , or heterocyclic ring;

A_7 is oxygen or $-ND_1$;


X_7 is a hydrogen or a halogen;

Y_7 is:

25



or X₇, A₇, and Y₇ taken together along with the carbons to which they are attached is:



5

wherein

R₆₁ is a hydrogen, a halogen, a lower alkyl group, -OD₁, or -NHC(O)O-lower alkyl;

R₆₂ is a hydrogen, a halogen, or a lower alkyl group; and

10 D₁ is as defined herein.

3. The compound of claim 2, wherein the compound is a benzimidazole, a quinoline, a pyrimidine, a thiadiazole, a sulfinylnicotinamide, a thienoimidazole, or a imidazopyridine.

4. The compound of claim 3, wherein the benzimidazole is
 15 omeprazole, lansoprazole, pantoprazole, rabeprazole, leminoprazole, timoprazole, tenatoprazole, disulprazole, esomeprazole, 2-(2-benzimidazolyl)-pyridine, a tricyclic imidazole, a thienopyridine benzimidazole, a fluoroalkoxy substituted benzimidazole, a dialkoxy benzimidazole, a N-substituted 2-(pyridylalkenesulfinyl) benzimidazole, a cycloheptenepyridine, a 5-pyrrolyl-2-
 20 pyridylmethylsulfinyl benzimidazole, a alkylsulfinyl benzimidazole, a fluoro-pyridylmethylsulfinyl benzimidazole, an imidazo[4,5-b]pyridine, RO 18-5362

or IY 81149; wherein the quinoline is a 4-amino-3-carbonyl quinoline, a 4-amino-3-acylnaphthyridine, a 4-aminoquinoline, a 4-amino-3-acylquinoline or a 3-butyryl-4-(2-methylphenylamino)-8-(2-hydroxyethoxy)quinoline; wherein the pyrimidine is a quinazoline, a tetrahydroisoquinolin-2-yl pyrimidine or YH 1885; wherein the thiadiazole is 3-substituted 1,2,4-thiadiazolo[4,5-a]benzimidazole or a 3-substituted imidazo[1,2-d]-thiadiazole; wherein the sulfinylnicotinamide is a 2-sulfinylnicotinamide; wherein the thienoimidazole is a pyridylsulfinylbenzimidazole, a pyridylsulfinyl thienoimidazole, a thienoimidazole-toluidine, a 4,5-dihydrooxazole, a thienoimidazole-toluidine or Hoe-731; wherein the imidazopyridine is a imidazo[1,2-a]pyridine, a pyrrolo[2,3-b]pyridine or a pharmaceutically acceptable salt thereof.

5 5. A composition comprising the compound of claim 2 and a pharmaceutically acceptable carrier.

15 6. The composition of claim 5, further comprising at least one of a nonsteroidal antiinflammatory drug, a selective COX-2 inhibitor, an antacid, a bismuth-containing reagent, and an acid-degradable antibacterial compound.

 7. A method for treating or preventing a gastrointestinal disorder, facilitating ulcer healing, or decreasing the recurrence of an ulcer in a patient in need thereof comprising administering to the patient a therapeutically effective amount of the composition of claim 5.

20 8. The method of claim 7, further comprising administering to the patient a therapeutically effective amount of an antacid.

 9. The method of claim 7, wherein the gastrointestinal disorder is an inflammatory bowel disease, Crohn's disease, irritable bowel syndrome, ulcerative colitis, a peptic ulcer, a stress ulcers, a bleeding peptic ulcer, a duodenal ulcer, infectious enteritis, colitis, diverticulitis, gastric hyperacidity, dyspepsia, gastroparesis, Zollinger-Ellison syndrome, gastroesophageal reflux disease, *Helicobacter Pylori* associated disease, short-bowel syndrome, or a hypersecretory state associated with systemic mastocytosis or basophilic leukemia and hyperhistaminemia.

30 10. A method for improving the gastroprotective properties, the anti-*Helicobacter pylori* properties, or the antacid properties of a proton pump

inhibitor comprising administering to a patient in need thereof a therapeutically effective amount of the composition of claim 5.

11. The method of claim 10, further comprising administering to the patient a therapeutically effective amount of a bismuth-containing reagent.

5 12. A method for decreasing or reversing gastrointestinal toxicity or facilitating ulcer healing resulting from administration of a nonsteroidal antiinflammatory drug and/or a selective COX-2 inhibitor to a patient comprising administering to a patient in need thereof a therapeutically effective amount of at least one composition of claim 5, and, optionally, at least one
10 nonsteroidal antiinflammatory drug and/or selective COX-2 inhibitor.

13. A method for treating *Helicobacter pylori* comprising administering to a patient in need thereof a therapeutically effective amount of at least one acid degradable antibacterial compound and at least one composition of claim 5.

14. A method for treating a viral infection comprising administering to
15 a patient in need thereof a therapeutically effective amount of the composition of claim 5.

15. The method of claim 14, wherein the viral infection is orthomyxoviridae, paramyxoviridae, picornaviridae, rhabdoviridae, coronavaridae, togaviridae, bunyaviridae, arenaviridae, reteroviridae,
20 adenoviridae, proxviridae, papovaviridae, herpesviridae, herpesviridae, herpes simplex viruses, cytomegalovirus, herpes varicella-zoster, Epstein-Barr, HHV6, HHV7, pseudorabies or rhinotracheitis.

16. A composition comprising at least one compound of claim 2 or a pharmaceutically acceptable salt thereof, and at least one compound that donates,
25 transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase.

17. The composition of claim 16 further comprising a pharmaceutically acceptable carrier.

30 18. The composition of claim 16, wherein the compound that donates, transfers, or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor or is a substrate for nitric oxide synthase is an S-nitrosothiol.

19. The composition of claim 18, wherein the S-nitrosothiol is S-nitroso-N-acetylcysteine, S-nitroso-captopril, S-nitroso-N-acetylpenicillamine, S-nitroso-homocysteine, S-nitroso-cysteine or S-nitroso-glutathione.

20. The composition of claim 16, wherein the S-nitrosothiol is:

- 5 (i) $\text{HS}(\text{C}(\text{R}_e)(\text{R}_f))_m\text{SNO}$;
 (ii) $\text{ONS}(\text{C}(\text{R}_e)(\text{R}_f))_m\text{R}_e$; and
 (iii) $\text{H}_2\text{N}-\text{CH}(\text{CO}_2\text{H})-(\text{CH}_2)_m-\text{C}(\text{O})\text{NH}-\text{CH}(\text{CH}_2\text{SNO})-\text{C}(\text{O})\text{NH}-\text{CH}_2-\text{CO}_2\text{H}$;
 wherein m is an integer from 2 to 20; R_e and R_f are each independently a
 hydrogen, an alkyl, a cycloalkoxy, a halogen, a hydroxy, an hydroxyalkyl, an
 10 alkoxyalkyl, an arylheterocyclic ring, an alkylaryl, a cycloalkylalkyl, a
 heterocyclicalkyl, an alkoxy, a haloalkoxy, an amino, an alkylamino, a
 dialkylamino, an arylamino, a diarylamino, an alkylarylamino, an
 alkoxyhaloalkyl, a haloalkoxy, a sulfonic acid, a sulfonic ester, an alkylsulfonic
 acid, an arylsulfonic acid, an arylalkoxy, an alkylthio, an arylthio, a cycloalkylthio,
 15 a cycloalkenyl, a cyano, an aminoalkyl, an aminoaryl, an aryl, an arylalkyl, an
 alkylaryl, a carboxamido, an alkylcarboxamido, an arylcarboxamido, an amidyl, a
 carboxyl, a carbamoyl, an alkylcarboxylic acid, an arylcarboxylic acid, an
 alkylcarbonyl, an arylcarbonyl, an ester, a carboxylic ester, an alkylcarboxylic ester,
 an arylcarboxylic ester, a haloalkoxy, a sulfonamido, an alkylsulfonamido, an
 20 arylsulfonamido, a sulfonic ester, a urea, a phosphoryl, a nitro, -T-Q, or
 $(\text{C}(\text{R}_e)(\text{R}_f))_k\text{-T-Q}$, or R_e and R_f taken together with the carbons to which they are
 attached form a carbonyl, a methanthial, a heterocyclic ring, a cycloalkyl group or
 a bridged cycloalkyl group; Q is -NO or -NO₂; and T is independently a covalent
 bond, a carbonyl, an oxygen, -S(O)_o- or -N(R_a)R_i-, wherein o is an integer from 0 to
 25 2, R_a is a lone pair of electrons, a hydrogen or an alkyl group; R_i is a hydrogen, an
 alkyl, an aryl, an alkylcarboxylic acid, an aryl carboxylic acid, an alkylcarboxylic
 ester, an arylcarboxylic ester, an alkylcarboxamido, an arylcarboxamido, an
 alkylaryl, an alkylsulfinyl, an alkylsulfonyl, an arylsulfinyl, an arylsulfonyl, a
 sulfonamido, a carboxamido, a carboxylic ester, an amino alkyl, an amino aryl,
 30 -CH₂-C(T-Q)(R_e)(R_f), or -(N₂O₂-)•M⁺, wherein M⁺ is an organic or inorganic
 cation; with the proviso that when R_i is -CH₂-C(T-Q)(R_e)(R_f) or -(N₂O₂-)•M⁺; then
 "-T-Q" can be a hydrogen, an alkyl group, an alkoxyalkyl group, an aminoalkyl
 group, a hydroxy group or an aryl group.

21. The composition of claim 16, wherein the compound that donates, transfers, or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase, is L-arginine, L-homoarginine, N-hydroxy-L-arginine, nitrosated L-arginine, nitrosylated L-arginine, nitrosated N-hydroxy-L-arginine, nitrosylated N-hydroxy-L-arginine, citrulline, ornithine, glutamine, lysine, polypeptides comprising at least one of these amino acids or inhibitors of the enzyme arginase.

22. The composition of claim 16, wherein the compound that donates, transfers, or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase is:

- (i) a compound that comprises at least one ON-O-, ON-N- or ON-C-group;
- (ii) a compound that comprises at least one O₂N-O-, O₂N-N-, O₂N-S- or -O₂N-C- group;
- (iii) a N-oxo-N-nitrosoamine having the formula: R¹R²-N(O-M⁺)-NO, wherein R¹ and R² are each independently a polypeptide, an amino acid, a sugar, an oligonucleotide, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted hydrocarbon, or a heterocyclic group, and M⁺ is an organic or inorganic cation.

23. The composition of claim 22, wherein the compound comprising at least one ON-O-, ON-N- or ON-C- group is an ON-O-polypeptide, an ON-N-polypeptide, an ON-C-polypeptide, an ON-O-amino acid, an ON-N-amino acid, an ON-C-amino acid, an ON-O-sugar, an ON-N-sugar, an ON-C-sugar, an ON-O-oligonucleotide, an ON-N-oligonucleotide, an ON-C-oligonucleotide, a straight or branched, saturated or unsaturated, substituted or unsubstituted, aliphatic or aromatic ON-O-hydrocarbon, a straight or branched, saturated or unsaturated, substituted or unsubstituted, aliphatic or aromatic ON-N-hydrocarbon, a straight or branched, saturated or unsaturated, substituted or unsubstituted, aliphatic or aromatic ON-C-hydrocarbon, an ON-O-heterocyclic compound, an ON-N-heterocyclic compound or a ON-C-heterocyclic compound.

24. The composition of claim 22, wherein compound comprising at least one O₂N-O-, O₂N-N-, O₂N-S- or O₂N-C- group is an O₂N-O-polypeptide, an

O₂N-N-polypeptide, an O₂N-S-polypeptide, an O₂N-C-polypeptide, an O₂N-O-amino acid, O₂N-N-amino acid, O₂N-S-amino acid, an O₂N-C-amino acid, an O₂N-O-sugar, an O₂N-N-sugar, O₂N-S-sugar, an O₂N-C-sugar, an O₂N-O-oligonucleotide, an O₂N-N-oligonucleotide, an O₂N-S-oligonucleotide, an O₂N-C-oligonucleotide, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted O₂N-O-hydrocarbon, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted O₂N-N-hydrocarbon, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted O₂N-S-hydrocarbon, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted O₂N-C-hydrocarbon, an O₂N-O-heterocyclic compound, an O₂N-N-heterocyclic compound, an O₂N-S-heterocyclic compound or an O₂N-C-heterocyclic compound.

25. The composition of claim 16, further comprising at least one of a nonsteroidal antiinflammatory drug, a selective COX-2 inhibitor, an antacid, a bismuth-containing reagent, and an acid-degradable antibacterial compound.

26. A method for treating or preventing a gastrointestinal disorder, facilitating ulcer healing, or decreasing the recurrence of an ulcer in a patient in need thereof comprising administering to the patient a therapeutically effective amount of the composition of claim 16.

27. The method of claim 26, further comprising administering to the patient a therapeutically effective amount of an antacid.

28. The method of claim 26, wherein the gastrointestinal disorder is an inflammatory bowel disease, Crohn's disease, irritable bowel syndrome, ulcerative colitis, a peptic ulcer, a stress ulcers, a bleeding peptic ulcer, a duodenal ulcer, infectious enteritis, colitis, diverticulitis, gastric hyperacidity, dyspepsia, gastroparesis, Zollinger-Ellison syndrome, gastroesophageal reflux disease, *Helicobacter Pylori* associated disease, short-bowel syndrome, or a hypersecretory state associated with systemic mastocytosis or basophilic leukemia and hyperhistaminemia.

29. A method for improving the gastroprotective properties, the anti-*Helicobacter pylori* properties, or the antacid properties of a proton pump inhibitor comprising administering to a patient in need thereof a therapeutically

effective amount of the composition of claim 16.

30. The method of claim 29, further comprising administering to the patient a therapeutically effective amount of a bismuth-containing reagent.

31. A method for decreasing or reversing gastrointestinal toxicity or facilitating ulcer healing resulting from administration of a nonsteroidal antiinflammatory drug and/or a selective COX-2 inhibitor to a patient comprising administering to a patient in need thereof a therapeutically effective amount of at least one composition of claim 16, and, optionally, at least one nonsteroidal antiinflammatory drug and/or selective COX-2 inhibitor.

32. A method for treating *Helicobacter pylori* comprising administering to a patient in need thereof a therapeutically effective amount of at least one acid degradable antibacterial compound and at least one composition of claim 16.

33. A method for treating a viral infection comprising administering to a patient in need thereof a therapeutically effective amount of the composition of claim 16.

34. The method of claim 33, wherein the viral infection is orthomyxoviridae, paramyxoviridae, picornaviridae, rhabdoviridae, coronaviridae, togaviridae, bunyaviridae, arenaviridae, reteroviridae, adenoviridae, proxviridae, papovaviridae, herpetoviridae, herpesviridae, herpes simplex viruses, cytomegalovirus, herpes varicella-zoster, Epstein-Barr, HHV6, HHV7, pseudorabies or rhinotracheitis.

35. A composition comprising at least one proton pump inhibitor compound or a pharmaceutically acceptable salt thereof, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase.

36. The composition of claim 35, wherein the at least one proton pump inhibitor compound is a benzimidazole, a quinoline, a pyrimidine, a thiadiazole, a sulfinylnicotinamide, a thienoimidazole, or a imidazopyridine.

37. The compound of claim 36, wherein the benzimidazole is omeprazole, lansoprazole, pantoprazole, rabeprazole, leminoprazole, timoprazole, tenatoprazole, disulprazole, esomeprazole, 2-(2-benzimidazolyl)-pyridine, a tricyclic imidazole, a thienopyridine benzimidazole, a fluoroalkoxy

substituted benzimidazole, a dialkoxy benzimidazole, a N-substituted 2-(pyridylalkenesulfinyl) benzimidazole, a cycloheptenepyrindine, a 5-pyrrolyl-2-pyridylmethylsulfinyl benzimidazole, a alkylsulfinyl benzimidazole, a fluoro-pyridylmethylsulfinyl benzimidazole, an imidazo[4,5-b]pyridine, RO 18-5362 or IY 81149; wherein the quinoline is a 4-amino-3-carbonyl quinoline, a 4-amino-3-acylnaphthyridine, a 4-aminoquinoline, a 4-amino-3-acylquinoline or a 3-butyryl-4-(2-methylphenylamino)-8-(2-hydroxyethoxy)quinoline; wherein the pyrimidine is a quinazoline, a tetrahydroisoquinolin-2-yl pyrimidine or YH 1885; wherein the thiadiazole is 3-substituted 1,2,4-thiadiazolo[4,5-a] benzimidazole or a 3-substituted imidazo[1,2-d]-thiadiazole; wherein the sulfinylnicotinamide is a 2-sulfinylnicotinamide; wherein the thienoimidazole is a pyridylsulfinylbenzimidazole, a pyridylsulfinyl thienoimidazole, a thienoimidazole-toluidine, a 4,5-dihydrooxazole, a thienoimidazole-toluidine or Hoe-731; wherein the imidazopyridine is a imidazo[1,2-a]pyridine, a pyrrolo[2,3-b]pyridine or a pharmaceutically acceptable salt thereof.

38. The composition of claim 37 further comprising a pharmaceutically acceptable carrier.

39. The composition of claim 35, wherein the compound that donates, transfers, or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase is an S-nitrosothiol.

40. The composition of claim 39, wherein the S-nitrosothiol is S-nitroso-N-acetylcysteine, S-nitroso-captopril, S-nitroso-N-acetylpenicillamine, S-nitroso-homocysteine, S-nitroso-cysteine or S-nitroso-glutathione.

41. The composition of claim 39, wherein the S-nitrosothiol is:

(i) $\text{HS}(\text{C}(\text{R}_e)(\text{R}_f))_m\text{SNO}$;

(ii) $\text{ONS}(\text{C}(\text{R}_e)(\text{R}_f))_m\text{R}_e$; and

(iii) $\text{H}_2\text{N}-\text{CH}(\text{CO}_2\text{H})-(\text{CH}_2)_m-\text{C}(\text{O})\text{NH}-\text{CH}(\text{CH}_2\text{SNO})-\text{C}(\text{O})\text{NH}-\text{CH}_2-\text{CO}_2\text{H}$;

wherein m is an integer from 2 to 20; R_e and R_f are each independently a hydrogen, an alkyl, a cycloalkoxy, a halogen, a hydroxy, an hydroxyalkyl, an alkoxyalkyl, an arylheterocyclic ring, an alkylaryl, a cycloalkylalkyl, a heterocyclicalkyl, an alkoxy, a haloalkoxy, an amino, an alkylamino, a

dialkylamino, an arylamino, a diarylamino, an alkylarylamino, an alkoxyhaloalkyl, a haloalkoxy, a sulfonic acid, a sulfonic ester, an alkylsulfonic acid, an arylsulfonic acid, an arylalkoxy, an alkylthio, an arylthio, a cycloalkylthio, a cycloalkenyl, a cyano, an aminoalkyl, an aminoaryl, an aryl, an arylalkyl, an alkylaryl, a carboxamido, a alkylcarboxamido, an arylcarboxamido, an amidyl, a carboxyl, a carbamoyl, an alkylcarboxylic acid, an arylcarboxylic acid, an alkylcarbonyl, an arylcarbonyl, an ester, a carboxylic ester, an alkylcarboxylic ester, an arylcarboxylic ester, a haloalkoxy, a sulfonamido, an alkylsulfonamido, an arylsulfonamido, a sulfonic ester, a urea, a phosphoryl, a nitro, -T-Q, or $(C(R_e)(R_f))_k$ -T-Q, or R_e and R_f taken together with the carbons to which they are attached form a carbonyl, a methanthial, a heterocyclic ring, a cycloalkyl group or a bridged cycloalkyl group; Q is -NO or -NO₂; and T is independently a covalent bond, a carbonyl, an oxygen, -S(O)_o- or -N(R_a)R_i-, wherein o is an integer from 0 to 2, R_a is a lone pair of electrons, a hydrogen or an alkyl group; R_i is a hydrogen, an alkyl, an aryl, an alkylcarboxylic acid, an aryl carboxylic acid, an alkylcarboxylic ester, an arylcarboxylic ester, an alkylcarboxamido, an arylcarboxamido, an alkylaryl, an alkylsulfinyl, an alkylsulfonyl, an arylsulfinyl, an arylsulfonyl, a sulfonamido, a carboxamido, a carboxylic ester, an amino alkyl, an amino aryl, -CH₂-C(T-Q)(R_e)(R_f), or -(N₂O₂-)•M⁺, wherein M⁺ is an organic or inorganic cation; with the proviso that when R_i is -CH₂-C(T-Q)(R_e)(R_f) or -(N₂O₂-)•M⁺; then "-T-Q" can be a hydrogen, an alkyl group, an alkoxyalkyl group, an aminoalkyl group, a hydroxy group or an aryl group.

42. The composition of claim 35, wherein the compound that donates, transfers, or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase is L-arginine, L-homoarginine, N-hydroxy-L-arginine, nitrosated L-arginine, nitrosylated L-arginine, nitrosated N-hydroxy-L-arginine, nitrosylated N-hydroxy-L-arginine, citrulline, ornithine, glutamine, lysine, polypeptides comprising at least one of these amino acids or inhibitors of the enzyme arginase.

43. The composition of claim 35, wherein the compound that donates, transfers, or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase is:

(i) a compound that comprises at least one ON-O-, ON-N- or ON-C-group;

(ii) a compound that comprises at least one O₂N-O-, O₂N-N-, O₂N-S- or -O₂N-C- group;

5 (iii) a N-oxo-N-nitrosoamine having the formula: R¹R²-N(O-M⁺)-NO, wherein R¹ and R² are each independently a polypeptide, an amino acid, a sugar, an oligonucleotide, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted hydrocarbon, or a heterocyclic group, and M⁺ is an organic or inorganic cation.

10 44. The composition of claim 43, wherein the compound comprising at least one ON-O-, ON-N- or ON-C- group is an ON-O-polypeptide, an ON-N-polypeptide, an ON-C-polypeptide, an ON-O-amino acid, an ON-N-amino acid, an ON-C-amino acid, an ON-O-sugar, an ON-N-sugar, an ON-C-sugar, an ON-O-oligonucleotide, an ON-N-oligonucleotide, an ON-C-oligonucleotide, a straight or
15 branched, saturated or unsaturated, substituted or unsubstituted, aliphatic or aromatic ON-O-hydrocarbon, a straight or branched, saturated or unsaturated, substituted or unsubstituted, aliphatic or aromatic ON-N-hydrocarbon, a straight or branched, saturated or unsaturated, substituted or unsubstituted, aliphatic or aromatic ON-C-hydrocarbon, an ON-O-heterocyclic compound, an ON-N-
20 heterocyclic compound or a ON-C-heterocyclic compound.

45. The composition of claim 43, wherein compound comprising at least one O₂N-O-, O₂N-N-, O₂N-S- or O₂N-C- group is an O₂N-O-polypeptide, an O₂N-N-polypeptide, an O₂N-S-polypeptide, an O₂N-C-polypeptide, an O₂N-O-amino acid, O₂N-N-amino acid, O₂N-S-amino acid, an O₂N-C-amino acid, an
25 O₂N-O-sugar, an O₂N-N-sugar, O₂N-S-sugar, an O₂N-C-sugar, an O₂N-O-oligonucleotide, an O₂N-N-oligonucleotide, an O₂N-S-oligonucleotide, an O₂N-C-oligonucleotide, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted O₂N-O-hydrocarbon, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or
30 unsubstituted O₂N-N-hydrocarbon, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted O₂N-S-hydrocarbon, a straight or branched, saturated or unsaturated, aliphatic or aromatic, substituted or unsubstituted O₂N-C-hydrocarbon, an O₂N-O-

heterocyclic compound, an O₂N-N-heterocyclic compound, an O₂N-S-heterocyclic compound or an O₂N-C-heterocyclic compound.

46. The composition of claim 35, further comprising at least one of a nonsteroidal antiinflammatory drug, a selective COX-2 inhibitor, an antacid, a bismuth-containing reagent and an acid-degradable antibacterial compound.

47. A method for treating or preventing a gastrointestinal disorder, facilitating ulcer healing, or decreasing the recurrence of an ulcer in a patient in need thereof comprising administering to the patient a therapeutically effective amount of the composition of claim 35.

48. The method of claim 47, further comprising administering to the patient a therapeutically effective amount of an antacid.

49. The method of claim 47, wherein the gastrointestinal disorder is an inflammatory bowel disease, Crohn's disease, irritable bowel syndrome, ulcerative colitis, a peptic ulcer, a stress ulcers, a bleeding peptic ulcer, a duodenal ulcer, infectious enteritis, colitis, diverticulitis, gastric hyperacidity, dyspepsia, gastroparesis, Zollinger-Ellison syndrome, gastroesophageal reflux disease, *Helicobacter Pylori* associated disease, short-bowel syndrome, or a hypersecretory state associated with systemic mastocytosis or basophilic leukemia and hyperhistaminemia.

50. A method for improving the gastroprotective properties, the anti-*Helicobacter pylori* properties, or the antacid properties of a proton pump inhibitor comprising administering to a patient in need thereof a therapeutically effective amount of the composition of claim 35.

51. The method of claim 50, further comprising administering to the patient a therapeutically effective amount of a bismuth-containing reagent.

52. A method for decreasing or reversing gastrointestinal toxicity or facilitating ulcer healing resulting from administration of a nonsteroidal antiinflammatory drug and/or a selective COX-2 inhibitor to a patient comprising administering to a patient in need thereof a therapeutically effective amount of at least one composition of claim 35, and, optionally, at least one nonsteroidal antiinflammatory drug and/or selective COX-2 inhibitor.

53. A method for treating *Helicobacter pylori* comprising administering to a patient in need thereof a therapeutically effective amount of at least one acid

degradable antibacterial compound and at least one composition of claim 35.

54. A method for treating a viral infection comprising administering to a patient in need thereof a therapeutically effective amount of the composition of claim 35.

5 55. The method of claim 54, wherein the viral infection is orthomyxoviridae, paramyxoviridae, picornaviridae, rhabdoviridae, coronavaridae, togaviridae, bunyaviridae, arenaviridae, reteroviridae, adenoviridae, proxviridae, papovaviridae, herpesviridae, herpes simplex viruses, cytomegalovirus, herpes varicella-zoster, Epstein-Barr, HHV6, 10 HHV7, pseudorabies or rhinotracheitis.

56. A method for preventing or treating a gastrointestinal disorder, facilitating ulcer healing, or decreasing the recurrence of an ulcer in a patient in need thereof comprising administering to the patient a therapeutically effective amount of the composition of claim 5 or a pharmaceutically acceptable salt 15 thereof, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase.

57. The method of claim 56, further comprising administering at least one antacid.

20 58. The method of claim 56, wherein the gastrointestinal disorder is an inflammatory bowel disease, Crohn's disease, irritable bowel syndrome, ulcerative colitis, a peptic ulcer, a stress ulcers, a bleeding peptic ulcer, a duodenal ulcer, infectious enteritis, colitis, diverticulitis, gastric hyperacidity, dyspepsia, gastroparesis, Zollinger-Ellison syndrome, gastroesophageal reflux disease, 25 *Helicobacter Pylori* associated disease, short-bowel syndrome, or a hypersecretory state associated with systemic mastocytosis or basophilic leukemia and hyperhistaminemia.

59. A method for preventing or treating a gastrointestinal disorder, facilitating ulcer healing, or decreasing the recurrence of an ulcer in a patient in need thereof comprising administering to the patient a therapeutically effective 30 amount of at least one proton pump inhibitor or a pharmaceutically acceptable salt thereof, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-

derived relaxing factor, or is a substrate for nitric oxide synthase.

60. The method of claim 59, further comprising administering at least one antacid.

61. The method of claim 59, wherein the gastrointestinal disorder
5 wherein the gastrointestinal disorder is an inflammatory bowel disease, Crohn's disease, irritable bowel syndrome, ulcerative colitis, a peptic ulcer, a stress ulcers, a bleeding peptic ulcer, a duodenal ulcer, infectious enteritis, colitis, diverticulitis, gastric hyperacidity, dyspepsia, gastroparesis, Zollinger-Ellison syndrome, gastroesophageal reflux disease, *Helicobacter Pylori* associated disease,
10 short-bowel syndrome, or a hypersecretory state associated with systemic mastocytosis or basophilic leukemia and hyperhistaminemia.

62. A method for improving the gastroprotective properties, the anti-*Helicobacter* properties or the antacid properties of a proton pump inhibitor compound comprising administering to a patient in need thereof a
15 therapeutically effective amount of a bismuth complex comprising at least one composition of claim 5.

63. A method for improving the gastroprotective properties, the anti-*Helicobacter* properties or the antacid properties of a proton pump inhibitor compound comprising administering to a patient in need thereof a
20 therapeutically effective amount of a bismuth complex comprising at least one composition of claim 16.

64. A method for improving the gastroprotective properties, the anti-*Helicobacter* properties or the antacid properties of a proton pump inhibitor compound comprising administering to a patient in need thereof a
25 therapeutically effective amount of a bismuth complex comprising at least one composition of claim 35.

65. A method for decreasing or reversing gastrointestinal toxicity or facilitating ulcer healing resulting from administration of a nonsteroidal antiinflammatory drug and/or a selective COX-2 inhibitor to a patient
30 comprising administering to a patient in need thereof a therapeutically effective amount of at least one composition of claim 5, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate

for nitric oxide synthase, and, optionally, at least one nonsteroidal antiinflammatory drug and/or selective COX-2 inhibitor.

66. A method for decreasing or reversing gastrointestinal toxicity or facilitating ulcer healing resulting from administration of a nonsteroidal antiinflammatory drug and/or a selective COX-2 inhibitor to a patient comprising administering to a patient in need thereof a therapeutically effective amount of at least one proton pump inhibitor compound, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase, and, optionally, at least one nonsteroidal antiinflammatory drug and/or selective COX-2 inhibitor.

67. A method for treating *Helicobacter pylori* comprising administering to a patient in need thereof a therapeutically effective amount of at least one acid degradable antibacterial compound, at least one composition of claim 5 or a pharmaceutically acceptable salt thereof, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase.

68. A method for treating *Helicobacter pylori* comprising administering to a patient in need thereof a therapeutically effective amount of at least one acid degradable antibacterial compound, at least one proton pump inhibitor compound or a pharmaceutically acceptable salt thereof, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase.

69. A method for treating a viral infection comprising administering to a patient in need thereof a therapeutically effective amount of at least one composition of claim 5 and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase.

70. The method of claim 69, wherein the viral infection is orthomyxoviridae, paramyxoviridae, picornaviridae, rhabdoviridae, coronaviridae, togaviridae, bunyaviridae, arenaviridae, reteroviridae,

adenoviridae, proxviridae, papovaviridae, herpesviridae, herpes simplex viruses, cytomegalovirus, herpes varicella-zoster, Epstein-Barr, HHV6, HHV7, pseudorabies or rhinotracheitis.

5 71. A method for treating a viral infection comprising administering to a patient in need thereof a therapeutically effective amount of at least one proton pump inhibitor compound or a pharmaceutically acceptable salt thereof, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase.

10 72. The method of claim 71, wherein the viral infection is orthomyxoviridae, paramyxoviridae, picornaviridae, rhabdoviridae, coronaviridae, togaviridae, bunyaviridae, arenaviridae, reteroviridae, adenoviridae, proxviridae, papovaviridae, herpesviridae, herpes simplex viruses, cytomegalovirus, herpes varicella-zoster, Epstein-Barr, HHV6, 15 HHV7, pseudorabies or rhinotracheitis.

73. A kit comprising at least one compound of claim 2 or a pharmaceutically acceptable salt thereof, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide 20 synthase.

74. The kit of claim 73, wherein the compound of claim 2 or a pharmaceutically acceptable salt thereof, and the at least one compound that donates, transfers or releases nitric oxide, induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric 25 oxide synthase are separate components in the kit or are in the form of a composition in the kit.

75. The kit of claim 73, further comprising at least one of a nonsteroidal antiinflammatory drug, a selective COX-2 inhibitor, an antacid, a bismuth-containing reagent and an acid-degradable antibacterial compound.


30 76. A kit comprising at least one proton pump inhibitor compound or a pharmaceutically acceptable salt thereof, and at least one compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide

synthase.

5 77. The kit of claim 76, wherein the proton pump inhibitor compound or a pharmaceutically acceptable salt thereof, and the compound that donates, transfers or releases nitric oxide, or induces the production of endogenous nitric oxide or endothelium-derived relaxing factor, or is a substrate for nitric oxide synthase are separate components in the kit or are in the form of a composition in the kit.

10 78. The kit of claim 76, further comprising at least one of a nonsteroidal antiinflammatory drug, a selective COX-2 inhibitor, an antacid, a bismuth-containing reagent and an acid-degradable antibacterial compound.

Figure 1



INTERNATIONAL SEARCH REPORT

Int. l. application No.
PCT/US00/02524

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/437, 31/4184, 31/4439 ; C07D 471/04, 401/12, 235/28
US CL : 514/303, 338, 395 ; 546/118, 273.7 ; 548/307.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/303, 338, 395 ; 546/118, 273.7 ; 548/307.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE, APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,629,305 A (EEK et al) 13 May 1997 (13.05.97), claims 1-19.	1-78 (in part)
A	US 5,599,794 A (EEK et al) 04 February 1997 (04.02.97), claims 1-2.	1-78 (in part)

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

25 APRIL 2000

Date of mailing of the international search report

05 JUL 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CHANA AULAKH

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/02524

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-78(In Part)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/02524

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

- I. Claims 1-78 (in part) directed to, A proton pump inhibitor compound represented by compounds of formula (I), classified in class 546, subclass 118+.
- II. Claims 1-78 (in part) directed to, A proton pump inhibitor compound represented by compounds of formula (II), classified in class 546, subclass 152+.
- III. Claims 1-78 (in part) directed to, A proton pump inhibitor compound represented by compounds of formula (III), classified in class 544, subclass 242+.
- IV. Claims 1-78 (in part) directed to, A proton pump inhibitor compound represented by compounds of formula (IV), classified in class 548, subclass 126+.
- V. Claims 1-78 (in part) directed to, A proton pump inhibitor compound represented by compounds of formula (V), classified in class 546, subclass 255+.
- VI. Claims 1-78 (in part) directed to, A proton pump inhibitor compound represented by compounds of formula (VI), classified in class 548, subclass 303.1+.
- VII. Claims 1-78 (in part) directed to, A proton pump inhibitor compound represented by compounds of formula (VII), classified in class 546, subclass 121+.

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons:
There is no common core Which in the Markush Practice, is a significant structural element shared by all of the alternatives; see PCT Administrative Instructions Annex B Part I (f) (i) (B) (1).